



SHELF LIST



0 0301 0029681 0

# NORMENTAFELN

ZUR

## ENTWICKLUNGSGESCHICHTE DER WIRBELTIERE.

IN VERBINDUNG MIT

Dr. BULL-Glasgow, Dr. BOCKL-Haber, HEALD, Prof. Dr. BRUCHER-Brüssel, Prof. Dr. B. DEAN-Columbia-University, New York, U. S. A., Dr. H. FUCHS-Strassburg, Dr. GILLESPIE-Strassburg, Prof. Dr. G. GÜNTHER, Prof. Dr. B. HENSENBERG-Gießen, Prof. Dr. HERRMANN-Utrecht, Prof. Dr. GEORGE M. KEEL-Glasgow, Prof. Dr. LAURIE, Prof. Dr. TH. KILMAYER-Breslau, Prof. Dr. LA ROSCHER, Prof. Dr. P. M. MAYER-Gießen, Dr. NIELSEN-Copenhagen, Prof. Dr. C. S. MINOT-Boston, U. S. A., Prof. Dr. NICOLY-Paris, Prof. Dr. PETER-Göteborg, Prof. R. ROBERTS-Van Arden, U. S. A., Dr. SAKUMA-Tokio, Japan, Dr. SEVERANCE-Harvard Medical School, Boston, U. S. A., Prof. Dr. SIMON-Paul-Ludwig-Institut, München, Prof. Dr. SOROKIN-Würzburg, Prof. Dr. SOROKIN-Louisiana, Prof. Dr. TAYLOR-Wien, Dr. TAYLOR-Philadelphia, U. S. A., Prof. Dr. TOURNAY-France, Dr. VON KELL-Paris, Prof. WILMANS-Chicago, U. S. A.

HERAUSGEGEBEN VON

**PROF. DR. F. KEIBEL**, LL. D. (HARVARD),  
FREIBURG I. BR.

---

**ELFTES HEFT.**

**NORMAL PLATES OF THE DEVELOPMENT OF NECTURUS MACULOSUS.**

BY

**ALBERT C. EYCLESHYMER and JAMES M. WILSON.**

LEO'S CASE, ILL. — GILL, WICH.

WITH 3 PLATES



**JENA,**

VERLAG VON GUSTAV FISCHER

1915.

# Normentafeln zur Entwicklungsgeschichte der Wirbeltiere.

Herausgegeben von  
**Prof. Dr. F. Keibel.**

(Fortsetzung)

- VI Normentafel zur Entwicklungsgeschichte des Schweines (*Sus scrofa domestica*). Herausgegeben von Prof. Dr. F. Keibel. Mit 3 lithogr. Tafeln. 1900. Preis 20 Mark.
- VII Normentafel zur Entwicklungsgeschichte des Huhnes (*Gallus domesticus*). Herausgegeben von Prof. Dr. F. Keibel. Mit 3 lithogr. Tafeln. 1900. Preis 20 Mark.
- VIII Normentafel zur Entwicklungsgeschichte des *Certodus forsteri*. Herausgegeben von Prof. Dr. Rich. Semper. Mit 3 lithogr. Tafeln. 1900. Preis 20 Mark.
- IX Normentafel zur Entwicklungsgeschichte der Zauneidechse (*Lacerta agilis*). Herausgegeben von Karl Peter. Mit 3 lithogr. Tafeln und 14 Figuren im Text. Preis 25 Mark.
- X Normal Plates of the development of the Rabbit (*Lepus europaeus* L.). By Charles S. Minot and Edwing Taylor. Harvard Medical School Boston, Mass. With 5 plates and 14 figures in the text. 1900. Preis 20 Mark.
- XI Normentafel zur Entwicklungsgeschichte des Rehes (*Cervus Capreolus*). Von Dr. Tsunejiro Sakurai. 1. Teil. Japan. Herausgegeben von Prof. Dr. F. Keibel. Mit 3 lithogr. Tafeln und 14 Figuren im Text. Preis 20 Mark.
- XII Normentafeln zur Entwicklungsgeschichte des Koholduaki (*Tarsius spectrum* und des Plumplori (*Nycticebus tardigræus*). Von A. W. Hübner, Friedrich und Franz Keibel. 1. Teil. München. Von Franz Keibel. Mit 3 Tafeln und 14 Textfiguren. 1907. Preis 20 Mark.
- XIII Normentafel zur Entwicklungsgeschichte des Menschen. Von Franz Keibel. 1. Teil. Leipzig. Von Carl Ullz. 2. Teil. Stockholm. Herausgegeben von Prof. Broman-Sund, Prof. Hammar-Upsala und Prof. Tandler-Wien. Mit 3 Tafeln und 14 Figuren im Text. 1905. Preis 30 Mark.
- XIV Normentafeln zur Entwicklungsgeschichte des Kiebitzes (*Vauellus cristatus* Meyer). Von Prof. O. Grosser, Wien. 1. Teil. Prof. O. Tandler, Wien. Mit 3 Tafeln. 1900. Preis 7 Mark 50 Pf.
- XV Normal Plates of the Development of *Lepidosiren Paradoxa* and *Protopterus aeneus* by J. Graham Kerr. University of Glasgow. With 1 Figure in the text. 1900. Preis 20 Mark.

## Fauna Arctica.

Eine Zusammenstellung der arktischen Tierformen mit besonderer Berücksichtigung des Spitzbergen-Gebietes auf Grund der Ergebnisse der Deutschen Expedition in das Nördliche Eismeer im Jahre 1898.

Unter Mitwirkung zahlreicher Fachgenossen  
herausgegeben von

† Fritz Römer und † Fritz Schaudinn.

nach ihrem Tode fortgesetzt von

**Dr. August Brauer**

in Berlin.

Bestehen des Werkes:

- Band I.** Die Insekten. Herausgegeben von Fritz Römer und Fritz Schaudinn. 1900. Preis 20 Mark.  
F. Römer: Die Dipteren. F. Schaudinn: Einführung. Ph. d. S. Wiedemann: Retschricht. Mit 2 Karten, 12 Abbild. im Text.  
E. E. Schulze: Die Hymenopteren. Mit 1 Tafel. J. Thiele: Die Neuropteren (*Chilopterus nev. spec.*). Mit 1 Tafel.  
O. v. Linstow: Die Coleopteren. Mit 2 Tafeln. H. Ludwig: Arktische und nördliche Heteropteren. W. Küken-  
thal: Die Lepidopteren von Grönland. C. Schaeffer: Die arktischen und subarktischen Coleopteren.  
J. A. Grieg: Die Spinnentiere von Grönland. Mit 10 Figuren. W. Weltner: Die Crustaceen der Arktis. Mit  
1 Tafel. F. Bolle: Die Crustaceen der Arktis. Mit 1 Tafel. F. Bolle: Die Crustaceen der Arktis. Mit 1 Tafel.  
H. Lohmann: Die Crustaceen der Arktis. Mit 1 Tafel. W. May: Die Crustaceen der arktischen und sub-  
arktischen Fauna. Mit 10 Figuren. C. Zimmer: Die arktischen Crustaceen. Mit 6 Figuren.  
H. Ludwig: Die Crustaceen der Arktis. O. Bidentkap: Die Bryozoen. 1. Teil. Die Bryozoen von Spitzbergen und  
Finnland. 1900. Preis 20 Mark.
- Band II.** Die Crustaceen. Herausgegeben von Fritz Römer und Fritz Schaudinn. 1900. Preis 20 Mark.  
H. Pöge: Die Crustaceen der Arktis. F. Schaudinn: Die Crustaceen der Arktis. Geographische Verbreitung der Crustaceen.  
F. Möbius: Die Crustaceen der Arktis. F. Schaudinn: Die Crustaceen der Arktis. A. Pagenstecher: Die Crustaceen der Arktis.  
F. Römer: Die Crustaceen der Arktis. H. Kraus: Die Crustaceen der Arktis. H. Friese: Die Crustaceen der Arktis.  
G. Brödmann: Die Crustaceen der Arktis. Al. Mrázek: Arktische Crustaceen. Mit  
1 Tafel. 1900. Preis 20 Mark.
- Band III.** Die Crustaceen. Herausgegeben von Fritz Römer und Fritz Schaudinn. 1900. Preis 20 Mark.  
F. Römer: Die Crustaceen der Arktis. F. Schaudinn: Die Crustaceen der Arktis. C. Graf Attems:  
Die Crustaceen der Arktis. O. Bidentkap: Die Crustaceen der Arktis. R. Hartmeyer: Die Crustaceen der Arktis.  
F. Römer: Die Crustaceen der Arktis. C. Zimmer: Die Crustaceen der Arktis. H. H. Gran: Die  
Crustaceen der Arktis. Mit 1 Tafel. 1900. Preis 20 Mark.
- Band IV.** Die Crustaceen. Herausgegeben von Fritz Römer und Fritz Schaudinn. 1900. Preis 20 Mark.  
F. Römer: Die Crustaceen der Arktis. F. Schaudinn: Die Crustaceen der Arktis. H. Schalow:  
Die Crustaceen der Arktis. O. Bidentkap: Die Crustaceen der Arktis. H. Schalow: Die Crustaceen der Arktis.  
F. Römer: Die Crustaceen der Arktis. E. Doderlein: Arktische Crustaceen. J. Mersen-  
hagen: Die Crustaceen der Arktis. Embrik Strand: Die Crustaceen der Arktis.  
O. Maas: Die Crustaceen der Arktis. 1900. Preis 20 Mark.
- Band V.** Die Crustaceen. Herausgegeben von Fritz Römer und Fritz Schaudinn. 1900. Preis 20 Mark.  
F. Römer: Die Crustaceen der Arktis. F. Schaudinn: Die Crustaceen der Arktis. C. Apstein: Die Crustaceen der Arktis.  
J. C. H. de Meijere: Die Crustaceen der Arktis. J. C. H. de Meijere: Die Crustaceen der Arktis.  
F. Römer: Die Crustaceen der Arktis. Rudolff Kowarzik: Die Crustaceen der Arktis.  
Hjalmar Broch: Die Crustaceen der Arktis. Rudolff v. Ritter Záhony: Die Crustaceen der Arktis.  
B. Poppra: Die Crustaceen der Arktis. 1900. Preis 20 Mark.

# NORMENTAFELN

ZUR

## ENTWICKLUNGSGESCHICHTE DER WIRBELTIERE.

IN VERBINDUNG MIT

Dr. BLES-Glasgow, Dr. BOEKE-Helder, Holland, Prof. Dr. BRACHET-Brüssel, Prof. Dr. B. DEAN-Columbia University, New York, U. S. A., Dr. H. FUCHS-Strassburg, Dr. GLAESNER-Strassburg, Prof. Dr. O. GROSSER-Prag, Prof. Dr. B. HENNEBERG-Giessen, Prof. Dr. HUBRECHT-Utrecht, Prof. J. GRAHAM KERR-Glasgow, Prof. Dr. KOPSCII-Berlin, Dr. THILO KRUMBACH-Breslau, Prof. Dr. LUBOSCH-Jena, Prof. Dr. P. MARTIN-Giessen, Dr. NIERSTRASZ-Utrecht, Prof. Dr. C. S. MINOT-Boston, U. S. A., Prof. Dr. NICOLAS-Paris, Prof. Dr. PETER-Greifswald, Prof. REIGHARD-Ann Arbor, U. S. A., Dr. SAKURAI-Fukuoka, Japan, Dr. SCAMMON-Harvard Medical School, Boston, U. S. A., Prof. Dr. SEMON-Prinz-Ludwigshöhe bei München, Prof. Dr. SOBOTTA-Würzburg, Prof. Dr. SOULIÉ-Toulouse, Prof. Dr. TANDLER-Wien, Dr. TAYLOR-Philadelphia, U. S. A., Prof. Dr. TOURNEUX, Toulouse, Dr. VOELKER-Prag, Prof. WHITMAN-Chicago, U. S. A.

HERAUSGEGEBEN VON

**PROF. DR. F. KEIBEL**, LL. D. (HARVARD),  
FREIBURG I. BR.

ELFTES HEFT.

NORMAL PLATES OF THE DEVELOPMENT OF NECTURUS MACULOSUS.

BY

**ALBERT C. EYCLESHYMER and JAMES M. WILSON.**

ST. LOUIS UNIVERSITY, ST. LOUIS MO., U. S. A.

WITH 3 PLATES.



**JENA,**  
VERLAG VON GUSTAV FISCHER.

1910.



---

Alle Rechte vorbehalten.

---

## P r e f a c e .

The preparation of the normal tables and plates on *Necturus* was begun several years ago by Professor C. O. WHITMAN but on account of unavoidable circumstances the work was delayed. It was later taken up by Professor EYCLESYMER upon the suggestion of both Professor WHITMAN and Professor KEIBEL. While Professor WHITMAN has not directly participated in the later work, he has furnished the senior author with material and information without which it would have been impossible to complete the work. In the completion of the work the senior author has been fortunate in having the able coöperation of Professor JAMES M. WILSON.



# Table of Contents.

	page
Description of illustrations . . . . .	1
Introduction to tables . . . . .	6
Tables . . . . .	7
Some variations in external structures . . . . .	28
Cleavage . . . . .	28
Gastrulation . . . . .	28
Closure of neural folds . . . . .	28
Appearance of optic vesicles . . . . .	28
Neuromeres (?) . . . . .	28
Somites . . . . .	29
Lateral curvature of body . . . . .	29
External gills . . . . .	29
Limbs . . . . .	29
Pigmentation . . . . .	29
Variations in internal structures . . . . .	30
Notochord . . . . .	30
Eye . . . . .	30
Ear . . . . .	30
Nose . . . . .	30
Epiphysis . . . . .	30
Paraphysis . . . . .	30
Hypophysis . . . . .	31
Liver . . . . .	31
Pancreas . . . . .	31
Spleen . . . . .	31
Thymus . . . . .	31
Trachea lungs . . . . .	31
Bibliography . . . . .	32
Introduction . . . . .	32
A. Alphabetical arrangement of titles according to authors . . . . .	32
B. Alphabetical arrangement of authors under topics . . . . .	48
Plates.	

### Description of Illustrations.

The series of eggs, embryos and larvae of *Necturus*, from which the following descriptions and the appended illustrations were made, were collected May 15th, 1903 and kept at a water temperature of 17°—18° C. The illustrations are copied from the original water colored pictures which were made by Mr. LEONARD H. WILDER, under the direction of the senior author. It should be emphasized that the ages, measurements and illustrations are all made from the living objects.

**Fig. 1.** (× 10.)

Side view of egg 1 day 4 hrs. after deposition. The first cleavage groove has reached the lower pole of the egg. Second grooves extend to level of the equator of the egg.

**Fig. 2.** (× 10.)

Side view of egg 1 day 8 hrs. after deposition. The second cleavage grooves have reached the equator. The grooves of the third cleavage pass in meridional planes, but have not yet reached the equator.

**Fig. 3.** (× 10.)

Side view of egg 1 day 12 hrs. old. Five cleavage grooves have reached lower pole, dividing lower hemisphere into six segments.

**Fig. 4.** (× 10.)

Side view of egg 1 day 16 hrs. old. The greater number of cleavage grooves pass in meridional planes, many are latitudinal and some nearly radial. The upper surface of the egg shows sixteen segments, the lower nine.

**Fig. 5.** (× 10.)

Side view of egg 1 day 20 hrs. old. The upper surface of the egg shows some fifty segments, the lower nine.

**Fig. 6.** (× 10.)

Side view of egg 2 days 2 hrs. old. The upper surface of the egg shows more than one hundred segments, the lower twelve.

**Fig. 7.** (× 10.)

Side view of egg 2 days 7 hrs. old. The upper surface of egg shows about two hundred cells. The lower portion is in about same stage as described in Fig. 6.



**Fig. 8.** (× 10.)

Side view of egg 2 days 12 hrs. old. The upper surface of egg shows some five hundred cells, the lower about forty.

**Fig. 9.** (× 10.)

Top view of egg 4 days 4 hrs. old. Segmentation cavity shows through thin translucent roof. Blastopore not present.

**Fig. 10.** (× 10.)

Bottom view of egg 6 days 16 hrs. old. Crescentic blastopore. Line of invagination sharply separates large yolk cells from small cells of blastodisc.

**Fig. 11.** (× 10.)

Dorso-lateral view of egg 10 days 10 hrs. old. Large circular blastopore; faint indication of embryonic anlage.

**Fig. 12.** (× 10.)

Side view of egg 10 days 16 hrs. old. Large circular blastopore. Anlage of mesial portion of embryo above dorsal lip of blastopore. Segmentation cavity faintly outlined.

**Fig. 13.** (× 10.)

Top view of egg 13 days 3 hrs. old. Small circular blastopore. Embryonic anlage triangular in outline; lateral boundaries indistinct. First appearance of neural groove. Roof of segmentation cavity thinner, making its boundaries distinct.

**Fig. 14.** (× 10.)

Top view of egg 14 days 4 hrs. old. Blastopore smaller, lateral margins of anterior portion of embryo bounded by short broad ridges which are the beginnings of the lateral portions of the neural fold. At anterior margin of embryo there is a transverse crescentic ridge which is beginning of transverse portion of neural fold. Neural groove deep but does not extend either to transverse portion of neural fold or to blastopore. Segmentation cavity crescentic.

**Fig. 15.** (× 10.)

Top view of egg 14 days 19 hrs. old. Blastopore much reduced, circular. The yolk plug is not visible in this egg. Lateral and transverse portions of neural fold united to form continuous fold around anterior portion of embryo. Lateral boundaries of posterior portion of embryo not defined. Neural groove not as long, nor as distinct as in preceding stage. Dark crescentic area in front of embryo is segmentation cavity.

**Fig. 16.** (× 10.)

Top view of egg 15 days 15 hrs. old. Blastopore small, circular; yolk plug visible. Neural fold prominent, its free ends extend nearly to blastopore. Neural groove deep and narrow at anterior end, broad and shallow at posterior end, fades out just in front of blastopore. A part of the segmentation cavity is still apparent in front of the embryo.

**Fig. 17.** (× 10.)

Top view of egg 16 days 6 hrs. old. Blastopore reduced to a very minute circular aperture. Neural plate narrower than in preceding stage. Neural fold prominent, its free ends coalescing at blastopore.

Neural groove extends to transverse portion of fold but does not reach blastopore. Segmentation cavity no longer visible in surface views.

**Fig. 18.** (× 10.)

Top view of embryo 17 days 2 hrs. old. Blastopore an elongated narrow aperture between ends of neural fold. Neural plate narrower than in preceding stage. The constricted portion represents in a general way the division between head and trunk. Neural fold most prominent in head region.

**Fig. 19.** (× 10.)

Top view of egg 17 days 17 hrs. old. Blastopore no longer visible. Neural plate narrowest posteriorly; broad in head region, showing boundary zone between head and trunk. Lateral portions of fold coalesced at posterior end of embryo. At anterior end of embryo a deep groove partially separates the two halves of the neural fold.

**Fig. 20.** (× 10.)

Top view of egg 18 days 13 hrs. old. Lateral portions of neural fold almost united except in head region where they are still widely separated. In the antero-lateral portions of the fold are slight evaginations which are the beginnings of the optic vesicles.

**Fig. 21.** (× 10.)

Top view of egg 18 days 15 hrs. old, 3 or 4 pairs of myotomes. Lateral portions of neural fold widely separated in head region, more closely approximated in anterior trunk region, coalesced in tail.

**Fig. 22.** (× 10.)

Dorso-lateral view of embryo 20 days 10 hrs. old, length 6 mm, 6 pairs of myotomes. Outline of body conforms to curvature of egg. Head end of embryo shows three longitudinal ridges; middle ridge lies slightly above level of lateral ridges. The middle one is common anlage of fore, mid and hind brain. The lateral ones are the common anlage of the optic vesicles and branchial arches. Anus formed.

**Fig. 23.** (× 10.)

Side view of embryo 21 days 2 hrs. old, length 7 mm, 10—12 pairs of myotomes. General outline of body conforms to curvature of egg. Head slightly raised above surface of yolk. Slight enlargement at end of tail. A distinct enlargement of anterior end of head shows optic vesicles; just posterior to this enlargement is the anlage of the branchial arches. Anus shows just below tip of tail.

**Fig. 24.** (× 10.)

Dorso-lateral view of embryo 22 days 17 hrs. old, length 8 mm, 16 - 18 pairs of myotomes. Embryo much curved laterally. Anterior half of head free from yolk. Caudal enlargement more prominent. Optic vesicles and mandibular arch well defined. The hyoid and first branchial arches are discernible; also the common anlage of the second and third branchial arches.

**Fig. 25.** (× 5.)

Side view of embryo 23 days 10 hrs. old, length 9 mm, 20—22 pairs of myotomes. General outline of the body straighter. Head free from yolk. Caudal enlargement becoming free. Optic vesicles and forebrain much larger. Mandibular, hyoid, first branchial, and common anlage of second and third branchial arches well defined. Otic vesicle visible above hyoid arch.

**Fig. 26.** (× 5.)

Side view of embryo 24 days 22 hrs. old, length 10 mm, 23—24 pairs of myotomes. General outline of body of embryo straighter, less curved laterally. Head and caudal extremities free from yolk. Yolk becoming oval. Optic vesicles prominent. Ear better defined. Olfactory pits present. The mandibular, hyoid and first branchial arches are distinct. The second and third branchial arches are not yet differentiated, a slight process on the first branchial indicates the beginning of the gill bar. The anlage of the heart is visible just beneath the arches.

**Fig. 27.** (× 5.)

Side view of embryo 26 days old, length 11 mm, 26—27 myotomes. General outline of body straighter than in preceding stage. Head projects some 3 mm beyond margin of yolk; tail projects 1.2 mm, is thinner laterally but broader dorso-ventrally. Eye, ear, nasal pits and mouth well defined. Maxillary process discernible. Mandibular arches longer, but ventral ends widely separated. Second and third branchial arches formed. Gill bars present on three branchial arches. Anterior limb buds indicated; faint anlage of posterior limb buds. Yolk pear-shaped. Heart prominent. First surface capillaries present although not indicated in figure.

**Fig. 28.** (× 5.)

Side view of embryo 30 days 8 hrs. old, length 13 mm, 30—31 myotomes. The trunk of the embryo is nearly straight. At level of the posterior gill there is a pronounced neck bend and at the level of the posterior limbs a striking downward bend of the tail. The epiphysis shows in surface views. The lens is discernible. The ear is still visible. The external nasal openings are sharply defined. The boundaries of the mouth are better outlined owing to the approximation of the ventral ends of the mandibular arches. The hyoid arch is becoming obscured. The gill bars are prominent on the three branchial arches. The anterior limb buds project dorsally about .5 mm above the surface of the body. The posterior limb buds are but slight elevations. The yolk is pear-shaped with its dorsal surface much flattened. The auricular and ventricular portions of the heart are apparent. The surface of the yolk is covered by a dense network of capillaries which for the most part convey blood antero-ventrally to the abdominal vein. Considerable pigment is present in the trunk region although but little has reached the outer portion of the dermis.

**Fig. 29.** (× 5.)

Side view of embryo 36 days 16 hrs. old, length 16 mm, 36—38 myotomes. In general outline the embryo shows a number of striking changes. The neck bend is not so pronounced. The tail bend is scarcely noticeable. There is a striking increase in dorso-ventral width of tail. The cerebral hemispheres are well defined. The eye is now prominent and the lens better defined. The ear is no longer visible in surface views. The mouth is well defined. The ends of the mandibular arches are closely approximated but not united. The hyoid and branchial arches are more obscure. Anlagen of gill filaments present on gill bars. Anterior limbs project dorsally. Posterior limbs are short ridges extending in horizontal plane. The yolk is elongated and reduced in diameter both dorso-ventrally and laterally. Surface blood vessels as in preceding stage, excepting that they are now apparent in the gill bars. The chromatophores are most numerous in the anterior and dorsal portions of the head.

**Fig. 30.** (× 5.)

Side view of embryo 40 days 20 hrs. old, length 18 mm, 44-46 myotomes. The outline of the body shows a marked ventral curvature of the trunk, less pronounced neck bend, and further increase in the dorso-ventral width of the tail. The eye is very prominent owing to the pigment in the retina. Ear not visible externally. Nasal openings very small. The mandibular arches have coalesced. The boundaries of the other arches are no longer discernible. Gill filaments well developed. Anterior limbs about 1 mm long project dorso-posteriorly. The yolk is elongated oval. Abdominal vein and branchial blood vessels prominent. Pigment present in dorsal portion of head, also along dorsal and lateral portions of trunk and tail. The yolk is unpigmented excepting along dorsal margin.

**Fig. 31.** (× 5.)

Side view of larva 49 days old, length 21 mm. General outline of body decidedly different. Head bend obliterated, slight upward curve in trunk. Tail broader. Eye more deeply pigmented. Gill bars very long, extending to level of end of anterior limb. From three to five lateral filaments on each gill bar. Anterior limbs project postero-ventrally; three digits formed. Posterior limbs directed caudad; no trace of digits. Yolk much elongated. Network of capillaries denser. Large lateral arteries, at level of upper margin of yolk, very prominent. Well defined longitudinal bands of pigment.

**Fig. 32.** (× 5.)

Side view of larva 61 days old, length 25 mm. General outline of body shows less dorsal curvature of trunk. Tail much longer in proportion to length of trunk and much broader dorso-ventrally. Gill bars longer, each possessing six to eight lateral filaments. Anterior and posterior limbs directed postero-ventrally. Anterior 3 mm long, posterior 2 mm long. Each limb shows four digits. The distribution of pigment is essentially similar to that observed in the 21 mm larva, the bands however are more sharply defined. Chromatophores in the gill bars and limbs and beginning to extend over the dorsal surface of the yolk.

**Fig. 33.** (× 5.)

Side view of larva 70 days 4 hrs. old, length 28 mm. The general outline of the body is slenderer than at any time preceding. The rapid absorption of the yolk has brought its ventral surface nearly to the level of the ventral surfaces of the head and tail. The gill bars curve dorsally and possess from ten to twelve pairs of lateral filaments. The tail is somewhat constricted at the level of the posterior limbs. The limbs and digits are better developed and are now used in locomotion. Pigmentation is denser than in 25 mm larva, but same general arrangement of bands prevails.

**Fig. 34.** (× 5.)

Side view of larva 97 days old, length 34 mm. In general outline the larva begins to resemble the adult. The yolk is well absorbed. The tail is very broad and now used as a powerful caudal fin in swimming. The gill bars project dorsally and have a large number of filaments. The legs project far below the ventral surface of the body. In color the same general pattern prevails as in the 28 mm larva.

There are some minor changes, the light band is broader and better defined, and extensions of pigment over the yolk have been so uneven that a number of irregular oval areas are left unpigmented, causing a mottled appearance in this region.

**Fig. 35.** (× 5.)

Side view of larva 126 days old, length 39 mm. The young *Necturus* now conforms in outline to the adult. In color however it is decidedly different.

### Introduction to Tables.

The material upon which the normal tables are based was collected in 1903. At the time the illustrations were made from the living material, several specimens of the same stages were fixed in various solutions. Among these 10% formalin caused the least distortion. This formalin-fixed material proved most satisfactory for work not involving cytological study. The material was stained *in toto* with haematoxylin, imbedded in paraffin and counterstained on the slide with weak picric acid, orange G or eosin. Unless otherwise stated the above fixation and staining have been used. Each of the stages designated in the following tables was sectioned in transverse, horizontal and sagittal planes. Besides these series many others have been consulted in which the material was fixed and stained by other methods. The total number of series at our disposal was upwards of 250.

# Tables.

---

Stage	Series	Length	Age	Blastomeres	Yolk
1	1		1 day 4 hrs.	2	
2	4		1 day 8 hrs.	6	
3	6		1 day 12 hrs.	12	Lower portion six surface grooves.
4	10		1 day 16 hrs.	20—24	Lower portion nine surface grooves; none reach center of egg.
5	12		2 days 2 hrs.	80—100	Lower portion twelve surface grooves; few reach center of egg.
6	18		2 days 7 hrs.	200—250	Lower surface as above excepting several grooves reach center of egg.
7	20		2 days 12 hrs.	500—600	Lower portion forty surface grooves; yolk much segmented.
8	25		6 days 16 hrs.		
9	28		10 days 16 hrs.		
10	35		13 days 3 hrs.		
11	38		14 days 4 hrs.		
12	40		14 days 10 hrs.		
13	48		15 days 10 hrs.		
14	55		16 days		
15	65		16 days 10 hrs.		
16	69		17 days		
17	73		17 days 17 hrs.		
18	85		18 days 15 hrs.		

Segmentation cavity	Blastopore. Anus	Archenteron. Enteron. Mesoderm. Chorda	Neurenteric canal	Nervous system. Optic vesicles. Somites	Stage
					1
					2
Present, large; roof single layer thick.					3
Large, roof in part two layers thick.					4
Segmentation cavity larger, roof in part two layers thick.					5
Segmentation cavity large, roof in part three layers thick.					6
Roof thinner, four layers around periphery, single layer in center.					7
Well formed.	Crescentic fissure on ventro-lateral portion of egg.	Archenteron just beginning.			8
Smaller.	Horse-shoe-shaped.	Archenteron extends over about 20°. Peristomal mesoblast present at dorsal lip of blastopore.			9
Smaller.	Circular.	Archenteron extends over about 45°. Dorsal wall posteriorly two layers. Peristomal mesoblast well defined in ventral lip of blastopore.			10
Quite small.	Small, circular, yolk plug externally visible.	Archenteron nearly complete; dorsal wall two layers anteriorly. Mesoblast around entire blastopore. Anlage of chorda.	Present.	Broad neural plate of thickened ectoblast; shallow neural groove.	11
Very small.	Very small, circular, yolk plug externally visible.	Archenteron very nearly complete; single layer in mid-longitudinal axis; this layer is beginning of chorda. Paraxial mesoblast co-extensive with chorda.	Present.	Low neural ridges present, deep neural groove.	12
Very small.	Elongated dorso-ventrally; yolk plug not externally visible.	Lateral walls of enteron three layers. Somatic and splanchnic layers well defined. Chorda well defined in head region.	Present, small.	Higher neural ridges, deep neural groove.	13
Very small.	Yolk plug not present.	Enteron single-layered dorsal wall; mesoblast extends ventrally over one-half of egg. Chorda narrower, thicker.	Tubular.	Prominent neural folds approaching each other, deep neural groove.	14
Not present (?).		Walls of enteron as above. Mesoblast extends to ventral portion of egg; well defined coelom. Chorda anteriorly a rod of cells.	Small canal.	High neural ridges approximating, but nowhere in contact.	15
Small, well defined.		Chorda oval or round in cross section throughout head and trunk regions, undifferentiated in tail.	Widely open.	Neural ridges meet in anterior head region. Beginning of optic vesicles.	16
(?)	Anus perforate.	Enteron as above. Mesoblast over ventral surface of egg. Chorda well defined in head and trunk regions.	Closed (?).	Neural ridges not closed in head region but coalesced in posterior trunk region and in tail. Optic vesicles better defined.	17
Present as large spaces among yolk cells.				Neural folds closed throughout entire length of embryo. Optic vesicles prominent, outer wall single layer of cells. Three or four somites.	18

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
19	100 Trans. 101 Trans.	6 mm	20 days 10 hrs.	Outline of body conforms to curvature of egg; extends over 160° of circumference of egg.	Externally 6 pairs of segments, those in anterior trunk region roughly triangular in cross sections. Myo-coele present. Muscle plate and cutis plate well defined. Clear zone at ends of muscle plate.				
19a		6 $\frac{1}{2}$ mm							
20	102 Trans. 103 Sag.	7 mm	21 days 2 hrs.	Body covers 180° of circumference of egg. Head end of embryo slightly elevated above surface of yolk. Distinct enlargement of anterior end shows optic vesicles. Tail end enlarged to form slight knob.	Externally 10 to 12 pairs of myotomes. In anterior trunk region muscle and cutis plate better defined, otherwise as in 6 mm stage.	Notochord anteriorly a solid rod of cells. Extends slightly in front of ear. Posteriorly not free from ectoderm and endoderm.	Neural tube closed throughout entire length; enlarged at anterior end. Fore brain, mid brain and hind brain indicated. Anlage of telencephalon. Fore brain nearly at right angles with mid brain; prominent, oval in transverse section; walls thin, cavity large. Mid brain long; oval in transverse section; canal narrow transversely. Optic stalk short, hollow. Chiasma well defined. Torus transversus small. Infundibulum moderate size. No plexus. No commissures. Anlage of trigeminal ganglion, also acusticus, facialis and vagus. Spinal cord long oval in section. Anlage of spinal ganglia. Caudally, notochord, dorsal wall of gut and neural tube fuse into common mass of cells.	Primary optic vesicles better defined; grown slightly backward; outer wall flattened and thickened forming anlage of retina; open widely into third ventricle. Anlage of lens as slight thickening of ectoderm. No mesoderm between lens and vesicle.	Otic vesicle just closed, spherical, lies between ectoderm and hind brain. Cells elongated in ventral part their inner ends free from yolk. Acustico-facialis ganglion lies ventro-cephalad of vesicle and is nearly as large as vesicle.
21	105 Trans.	8 mm	22 days 18 hrs.	Embryo curved laterally. Anterior half of head free from yolk. Optic vesicles well defined.	Externally 14 to 16 pairs myotomes.	Well defined throughout anterior and middle portion of body; extends slightly in front of level of ear. Anterior portion lies in contact with medullary tube. Long tract of mesenchyma between anterior end of chorda and hypophysis; fused with ectoderm and endoderm in posterior portion. Anlage of sheath. Indications of hypochorda.	Cranial flexure well marked. Divisions of brain better defined. Medulla longer than fore brain and mid brain together. In roof of fore brain, paraphysial arch, postvelar arch and epiphysial arch formed. Beginning of velum transversum. Trigeminal nerve forming branches. Roof of medulla a single layer of flattened cells; nerve fibers forming in Gasserian ganglion. Spinal cord oval in section. Canal wide, constricted at sides. Roof plate thinner than floor plate. Cells still contain large yolk granules. Spinal ganglia have grown down nearly to middle of side of cord.	Vesicle larger. Beginning of optic cup. Inner layer thinner; outer or retinal layer slightly concave. Optic stalk somewhat constricted.	Vesicle completely closed, slightly pear-shaped, smaller end dorsal. In contact with ectoderm.
22	106 Trans. 107 Sag. 110 Front	9 mm	23 days 10 hrs.	Head end elevated. Optic vesicles larger. Mandibular, hyoid, first branchial and common anlage of second and third branchial arches well defined. Caudal enlargement more pronounced. Embryo in contact with yolk over 140°. Curved laterally.	Externally 20 to 22 pairs of myotomes. In anterior trunk region, muscle plate nearer notochord. Clear zones at ends of plate wider, fibrillae present in myoblasts.	Extends from a point slightly in front of ear to tip of tail. Vacuolated anteriorly, not posteriorly. Separated from neural tube, fore gut and hypophysis by mesenchyma. Posteriorly notochord, neural tube and endoderm fused into common mass. Unsegmented. Hypochorda well defined.	Flexure showing in mid brain. Hemispheres well developed; walls thinner; ventricle very large. Beginning of epiphysis. Velum transversum deeper. Infundibulum forming. Optic chiasma well defined, also torus. Roof of mid brain thin, ventricle widest in dorsal portion. Auditory nerve well developed. Trigeminal ganglion has two branches, one running over eye, other anterior to first gill. Peripheral band of nerve fibers on ventral and lateral surface of medulla. Spinal cord elongated oval in transverse section. Canal wide. Roof plate thin, well defined. Ganglia larger. Undifferentiated mass of cells at caudal end of spinal cord.	Optic cup shallow. Optic stalk better defined; contains small tubular lumen which connects third ventricle of brain and cavity of optic vesicle. Lens as short dorsal evagination. Thin layer of mesenchyma connection with ectoderm and containing cavity.	Vesicle pear-shaped, smaller end dorsal, detached from ectoderm. Anlage of ductus endolymphaticus as short dorsal evagination. Thin layer of mesenchyma between vesicle and ectoderm, also between vesicle and hind brain.

Nose	Hypophysis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Stage
								19
							Single layer of long flat cells containing coarse yolk granules.	19a
Anlage of nasal organ as thickening of internal layer of ectoderm.	Anlage of hypophysis as a long wedge-shaped mass of cells the inner end of which is 2 or 3 layers thick, outer end a single layer continuous with deeper layer of ectoderm. No capsule. Lies some distance from anterior end of notochord. Wall closely applied to single-layered wall of infundibulum. Yolk granules less numerous than in underlying entoderm.	Position of mouth indicated by thickening of ectoderm.	Gut extends from level of posterior portion of eyes to a point slightly beyond posterior end of embryo. Distended anteriorly to form branchial chamber: short postanal gut.	Anlage of thyroid as median outgrowth in anterior portion of floor of branchial cavity, in close relation to hyomandibular arch. No external indications of gill arches.	Pronephric ducts present as two short straight tubes.	Anlage of heart in form of a rod of cells lying below the pharynx and between the ventral borders of the pericardial cavities.	Thickened greatly over head. Over body long flat cells interspersed with many large oval cells.	20
Thickening of internal layer of ectoderm more pronounced and more definitely circumscribed.			Anlage of liver.	Thyroid evagination deeper. Outgrowth longer. Mandibular hyoid and common Anlage of third, fourth and fifth arches visible externally.				21
Slight concavity in surface of ectoderm. Thickness of wall of nasal organ equal to that of wall of forebrain. Narrow layer of mesenchyma between nasal organ and forebrain. External layer of ectoderm can be traced only to margin of nasal organ.	Mass of cells close under infundibulum, 4-5 cell layers thick. Slight indications of cavity. Has lost connection with ectoderm.	Surface invagination a broad transverse groove.	Gut extends from anlage of mouth to junction of tail end of embryo with yolk; here it runs ventrad and extends somewhat further over the yolk than in the preceding stage. Extends laterally far beyond bounds of embryo at level of anterior portion of yolk. Diverticulum of gut just behind liver extends ventrally half way to ventral surface. Pharyngeal chamber large. Midgut narrow dorso-ventrally, wide laterally. No distinction between midgut and hindgut. Wall of pharynx and dorsal wall of midgut consist of single layer of columnar cells, heavily laden with yolk granules. Ventral wall of midgut irregular yolk cells. Liver pear-shaped evagination.	Mandibular, hyoid, first branchial and common Anlage of fourth and fifth arches visible externally.	Pronephric ducts coiled, open anteriorly. End posteriorly about 10" or 11" segment. End in mass of mesoderm in close proximity to entoderm.	Pericardial cavity large. Heart has become a tube. Endothelium distinct. Division into auricular and ventricular portion recognizable. Lateral blood vessels appearing, also branchial arches and sinus venosus.	Thickened over fore and mid brain.	22

Stage	Series	Length	Age	Body-form	Somites	Notochord	Nervous System	Eye	Ear
23	111 Trans. 112 Sag. 113 Front.	10 mm	24 days 22 hrs.	Head more elevated. Optic vesicles prominent. Embryo less curved ventrally and laterally. Tail end enlarged, becoming free from yolk. Yolk becoming oval in outline. Marked cephalic flexure.	Externally 23 to 24 pairs of myotomes. Muscle plate much wider laterally, extends further ventrally. Myocoel small.	Sheath well defined, a single layer of cells. Vacuolated throughout excepting a small portion at posterior end. Hypochorda well defined.	Flexure in mid brain more pronounced. Cerebral hemispheres better defined. Neuromeres in a single layer of cells; inner wall or physis indicated as thickening in epithelium. Spinal cord oval in transverse section. Layer of fibres well defined in ventrolateral portion of cord. Spinal ganglia better defined. Many nerve fibers beginning to grow out from ganglion cells.	Optic cup concave; outer pigment wall or lymphaticus better developed. Inner ends of cells becoming clear. Optic stalk narrower. Lens spherical, hollow, not completely detached from inner layer of ectoderm. Mesoderm extending between lens and wall of cup.	Vesicle pear-shaped. Ductus endolymphaticus better developed.
24	116 Trans. 117 Front.	11 mm	26 days	Outline of body straighter. Anterior limb buds indicated. Yolk pear-shaped. Heart prominent.	Externally 26 to 27 myotomes. In anterior trunk region muscle plate and cutis plate further differentiated; muscle plate extends ventrally to level of lower margin of notochord. Myocoel nearly obliterated. Septa between myotomes forming by ingrowing mesenchyma.	Extent as above. Separated from neural tube by mesenchyma throughout head and trunk. Separated from gut by mesenchyma and dorsal aorta. Hypochorda better defined.	Epiphysis better defined; paraphysis a slight evagination of both a single layer of cells. Velum transversum also a single layer of cells.	Optic cup more concave, cavity very small. Lens detached from ectoderm; inner wall thicker than outer. No cellular elements in cavity of lens. Mesoderm also extending between lens and optic cup.	Vesicle pear-shaped; lateral portion thick, ventral portion very thick. Ductus endolymphaticus being constricted.
25	118 Trans. 119 Sag.	12 mm	28 days 2 hrs.	Embryo less curved dorsally and laterally. Head end free to anlage of heart. Tail end projects further from yolk. Marked cephalic flexure. Ends of mandibular arches near median line but still separate. Yolk changed from oval form to pear-shaped, small end anterior. Capillaries appearing in yolk.	Externally 28 to 29 myotomes (?).	Extends to level of ear anteriorly. Separated from neural tube by mesenchyma throughout entire length except at caudal end. Separated from gut by mesenchyma and dorsal aorta.	Paraphysis a well marked narrow diverticulum. Spinal cord cells still arranged radially. Layer of fibers better defined.	Pigment wall of optic cup very thin, a single layer of flat cells. Optic stalk more constricted. Lens larger, nearly fills optic cup; cavity in lens larger. Mesoderm between lens and ectoderm. A few mesoderm cells in optic cup behind lens.	
26	120 Trans. 125 Sag. 126 Front.	13 mm	30 days 8 hrs.		Externally 30 to 31 myotomes.	Extends from otic vesicle to posterior end of spinal cord. Vacuolated throughout greater portion of its extent. Separated from neural tube and chiasma and torus. Commis-gut throughout entire extent. No mesenchyma between notochord and neural tube. Separated from gut by mesenchyma and dorsal aorta. Hypochorda not present (?).	Cervical flexure well marked. Diencephalon and mesencephalon nearly at right angles to medulla. Velum transversum much deeper. Infundibulum and hypophysis larger, also optic chiasma and torus. Commis-plexus. Diencephalon still projects in front of prosencephalon. Ganglion of trigeminal very large. Peripheral layer of fibers developed in ventral and lateral wall of mid brain. Beginning of plica rhombencephalica. Spinal cord oval in section, compressed laterally; canal a narrow slit. Spinal ganglia large, well developed.	Optic cup has outer margin in close contact with ectoderm. Cavity obliterated except at outer margin. Pigment wall and retinal layer as above. Lumen of optic stalk smaller. Choroideal groove deep. Walls of lens of uniform thickness.	Vesicle flattened on mesial surface; walls of saccular portion notably thicker than remaining portion. Mesoderm separates vesicle from ectoderm, also from hind brain. Ductus longer, constricted.

Nose	Hypophysis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
Disk-like depression on surface deeper. Nasal epithelium thicker, separated from fore-brain by thin layer of mesenchyma.				Gill bearing arches recognizable as divisions in a common enlargement from which they develop. Anlage of external gills as slight proliferations. Anlage of trachea as longitudinal groove in median line of thickened ventral wall of oesophagus. Gill clefts indicated by deep evaginations of pharyngeal and corresponding invagination of surface ectoderm. No ectodermal invagination between fourth and fifth arches.	Pronephric duct much coiled at anterior end; same caliber throughout; widely open anteriorly into coelom; posteriorly open into cloaca. No trace of mesonephros.	Heart somewhat twisted. Endothelium and mesothelium defined. Blood vessels present in anterior branchial arches. Truncus arteriosus formed. Beginnings of vitelline veins as irregular spaces on ventral surface of yolk. Anterior and posterior cardinal veins forming. Blood capillaries appearing over surface of yolk. No blood vessels in tail.			Anlage of anterior limbs visible.	23
Invagination deeper, cup-like. In free ends of cells forming bottom of cup, yolk is entirely absorbed leaving clear zone.	An oval loosely arranged mass of cells containing irregular cavities. Antero-ventral wall thinner than dorsal. Yolk not absorbed.			Gill bearing arches better developed. Ectodermal invagination between fourth and fifth arches. Tracheal groove deeper; wall a single layer of columnar cells.			Generally two layers. Cuticular margin well developed. Shows stratum corneum and stratum germinativum, each a single layer of cells. Many large oval cells (mucous, "Leydig" cells).			24
Nasal pit formed; organ oval in outline.		Epithelium of mouth shows as columnar layer of cells, although yolk is not absorbed in mouth cavity.	Anlage of gall bladder. Anlage of dorsal pancreas.	Gill bearing arches better defined. Gill bars prominent but no gill filaments. Pharyngeal portion of gill clefts extends two thirds distance to exterior, meeting ectodermal invaginations. Walls of clefts closely apposed.		Heart placed transversely. Large abdominal vein formed communicating with sinus venosus.			Anterior limbs more prominent. Posterior limb buds just discernible.	25
Nasal pit a short cylinder. Epithelium thicker. Clear zone at free ends of cells broader	An oval mass of cells closely applied to infundibulum. Contains cavity. Capsule beginning to be formed.			Thyroid extends backward to anterior wall of pericardium. Tracheal evagination deeper, tubular.	Pronephric tubules still widely open into coelom. Posterior portion narrower than anterior. No mesonephros.	Heart S-shaped, twisted. Auricular portion tubular. Ventricle constricted where it opens into conus. Conus and truncus not differentiated. Endothelium of ventricle and of pericardial cavity a single layer of flattened cells. Anterior and posterior cardinal veins formed. Large abdominal vein running through the liver. Beginning of portal circulation.	Cuticular margin thicker. Anlage of lateral line organs.			26



Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
27	127 Trans. 128 Sag. 130 Front.	14 mm	32 days 10 hrs	Trunk of embryo straight when viewed from side. Cephalic flexure still pronounced. Neck bend marked. Pronounced ventral bend in tail at level of posterior limbs. Lateral curvature pronounced. Yolk elongated, pear-shaped. Flat-topped on dorsal surface; large abdominal vein. Considerable pigment in trunk.	Externally 31 to 32 myotomes. In anterior trunk region myotomes much elongated dorso-ventrally, extend below level of notochord. Muscle plate covers one half distance between notochord and ectoderm. Cutis plate recognizable. Septa well differentiated, fibrillae fill one half of muscle cell.	Extent as above. Anterior end almost in contact with hypophysis, a small tract of mesenchyma intervening.	Epiphysis a small flattened vesicle; cavity in stalk obliterated. Posterior commissure indicated. Lumen in optic stalk very small. Dorsal wall of diencephalon a single layer of cells. Paraphysis and epiphysis both surrounded by mesenchyma. Plicarhomboencephalica a deep invagination. Infundibulum larger; ventral process extending caudad, posterior wall single layer of cells, closely applied to antero-ventral wall of medulla. Cord oval in outline in cross section, ventral half narrower than dorsal. Layer of fibers on outside of ventral half well defined. Yolk granules have nearly disappeared.	Margins of optic cup are in close contact with ectoderm. Wall of retina thicker at bottom of cup, thinner at margin. Pigment layer thinner, consisting of a single layer of flat cells. Optic stalks longer, lumen still present. Choroidal fissure present. Inner wall of lens thicker than outer, projecting into disk-like cavity.	Vesicle further flattened on mesial surface and elongated antero-posteriorly. Ductus elongated, narrowed and more constricted at base; extends well up on sides of medulla. External wall a single layer of flattened cells; mesial wall thinner.
28	131 Trans. 133 Sag. 134 Front.	15 mm	34 days 12 hrs		Shows 32-34 myotomes.	Anterior end close to hypophysis, some mesenchyma intervening; posterior reaches end of spinal cord. Anterior end slightly enlarged. Yolk absorbed in central portion. Caudal end still in contact with neural tube. Sheath better developed, not segmented. Hypochorda not present.	Velum transversum a single layer of cells. Beginning of choroid plexuses as two invaginations, one in front of paraphysis, other behind it. Posterior and anterior commissure appear. Infundibulum extended laterally. Commissura habenularis forming.	Margin of optic cup more constricted. Cavity of optic vesicle still present in margin of cup. Optic stalk smaller, longer, communicates with third ventricle by very narrow channel. Lens more completely fills optic cup; outer wall a single layer of cells; inner wall much thickened. Mesenchyma increasing between lens and ectoderm. A few scattering mesenchyma cells in optic cup. Choroidal fissure nearly closed.	Beginnings of lagena and semi-circular canals.
29	135 Trans. 138 Sag. 139 Front.	16 mm	36 days 16 hrs.	Trunk and tail straight line. Neck and cephalic flexures less pronounced. Tail greatly widened dorso-ventrally at level of posterior limbs. Yolk much elongated, ventral surface convex, dorsal slightly concave in profile view. Mandibular arches strongly developed, ventral ends enlarged, not yet united. Pigment present in trunk and dorsal portion of head. Anterior limb buds prominent, posterior slight elevations.	Horizontal section shows 36-38 myotomes. In anterior trunk region muscle plate extended ventrally to level of pronephric ducts. Cutis plate obscure.	Extends almost to hypophysis, some mesenchyma intervening. Knobbed at anterior end. Greatest diameter at level of anterior margin of yolk. Vacuolated throughout except near ends. In head and trunk considerable mesenchyma between notochord and neural tube. Sheath better developed. Mesenchymal thickenings in sheath indicate beginnings of intervertebral disks.	Walls of telencephalon differentiated into three layers, an outer fibrous layer, a middle layer of loose cells, and an inner layer of more compact cells. Chiasma contains few fibers. Paraphysis widely open, pear-shaped, some lateral diverticula. Anterior and posterior choroid plexuses formed. Posterior larger and more convoluted. Anterior and posterior commissures present. Anlage of commissura habenularis. Recessus lateralis small; dorsal, lateral and median longitudinal zones well defined. Tuberculum posterius formed; contains commissure	Optic cup more constricted at margin coming in contact with lens. Retina much thickened except at margin where it is abruptly thinned indicating ora serrata. Pigment beginning to be formed in retina. Clear zone in retina next to vitreous. Indications of eye muscles.	

Nose	Hypo-physis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urogenital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
Nasal tube curves dorso-caudally. Nasal organ an elongated oval. Becoming separated from superficial ectoderm by ingrowing mesenchyma. Condensed layer of mesenchyma around organ indicates beginning of fibrous capsule.			Dorsal pancreas further evaginated. Anlage of spleen as a small mass of mesenchymal cells in dorsal mesentery of stomach.	Gill-bearing arches less prominent. Gill bars better developed. Gill filaments recognizable as slight buds. Third and fourth clefts perforate. Tracheal evagination deeper, distal end dilated, becoming pear-shaped. Wall a single layer of cells.	Pronephro- stomic openings constricted.	Branchial arches all contain blood vessels.		Skull showsanlage of trabeculae in condensed mesenchyma.	Anterior limbs project dorsally. Posterior limb buds more prominent.	27
	Entirely free from foregut. Lies close against infundibulum. Surrounded by connective tissue capsule. Yolk absorbed. No trace of division into lobes. No cavity.	Sharply defined groove indicates position of mouth. Roof and floor of oral cavity clearly defined but cavity filled with yolk-bearing cells.	Walls of pharynx consist of one or two layers of flattened cells excepting posteriorly where still columnar. Oesophagus open. Stomach spindle-shaped, its walls composed of long columnar cells filled with yolk granules; surrounded by two or three layers of densely packed mesenchymal cells representing the beginnings of its muscular walls. Midgut small, dorsal wall fairly well defined, ventral wall irregular yolk cells. Cloaca a wide chamber, yolk nearly absorbed in its dorsal wall. Liver further extended laterally, consists of 15 to 20 hepatic cords separated by sinusoids. Cells heavily laden with yolk. Gall bladder small but well defined. Cystic duct and hepatic duct open in common into the wide ductus choledochus and this opens widely into the intestine.	Posterior portion of thyroid shows beginning of division into two parts. Tracheal diverticulum pear-shaped lying in mesenchyma in median line below oesophagus; cells columnar; yolk being rapidly absorbed. Just anterior to tracheal diverticulum marked thickening in ventral wall of gut.	Pronephric tubules lie opposite 2" or 3" somites. Mesonephros begins in 9" segment from front and extends to 17". Best developed anteriorly. Posteriorly not metameric.	Partition separating auricles nearly complete. Truncus arteriosus larger. Primitive aortae large. Hepatic vein very large. Blood vessels present in anterior limb buds.	Pigment appearing. Anlage of basement membrane. Sense organs developing.	Anlage of trabeculae, MECKEL'S cartilage palatoquadrate, basal plate, hyoid arch, 1", 2" and 3" branchial arches, first basibranchial and faint indications of second basibranchial, also of occipital arches, all in condensed mesenchyma. Mandibular bars separated from each other ventrally also separated from basi-cranial plates. Quadrate closely applied to MECKEL'S cartilage.		28
Nasal tube deeper. Nasal organ elongated oval. Condensation of mesenchyma around periphery of organ more marked.				Gill bars longer. Gill filaments beginning. Second cleft between hyoid and first branchial perforate, third cleft perforate, fourth perforate, fifth not perforate. Anlage of thymus as proliferations of epithelium of dorsal portions of gill arches. Tracheal evagination longer more dilated at distal end. Anlagen of lungs as lateral diverticula of distal end of trachea. Dorsal wall of trachea thicker than ventral.		Auricle small and nearly spherical; lies just dorso-cephalad from ventricle. Ventricle turned to the right. Constriction indicates valves between auricle and ventricle. Sinus venosus very large; it receives ductus Cuvieri, posterior cardinals and pronephric branch; also vein formed by internal and external jugulars. Vitelline vein very large; enters liver. Abdominal vein large, passes through liver, joins hepatic vein and other veins to form hepatic portal system, then enters sinus venosus. Large vessel on either side. Small vessels in posterior limbs. Vessels around nose and eye; also in pia mater. Dorsal aorta formed. Also subclavian and brachial arteries.	Cartilage appearing around outer portion of otic capsule.	Anterior limbs better developed; directed dorsally. Posterior limb buds slight elevations; long axis in horizontal plane.	29	

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
30	140 Trans. 144 Sag. 145 Front.	17 mm	38 days		Horizontal section shows 40-42 myotomes.	Yolk not absorbed in anterior end. Vascularized through-out. Mesenchyma between anterior end and hypophysis. Posteriorly rests against neural tube. Not segmented. Beginnings of neural arches in cartilages.	Plexuses much less marked. Anterior end of brain formed by hemispheres. Paraphysis well developed; plexus extends into lateral ventricles and diencephalon. Lamina terminalis thick. Anterior commissure well defined, also commissura habenularis. Lamina cerebellaris recognizable. Ganglion of trigeminal divided into two parts.	Optic cup flattened laterally, surrounded except on outer surface, by condensed mesenchyma. Retina divided into two layers; contains some pigment. Anlage of iris smaller, cavity very small. Lens larger, spherical. Inner wall fills cavity except narrow cleft. Anterior Spinal cord rounder; canal wall a single layer of shorter dorso-ventrally. Ganglia very large; nerve roots well developed; nerve fibers appearing in ventral nerve roots. Anlage of pia mater.	Ear more extended antero-posteriorly. Utriculo-saccular partition beginning. Ductus endolymphaticus extends over lateral margins of medulla toward median line; dilated at distal end to form saccus. Anlagen of lagenae and semicircular canals. Periotic capsule forming in condensed mesenchyma.
31	147 Trans. 156 Sag. 157 Front.	18 mm	40 days 20 hrs.	Trunk and tail slightly concave in profile. Cervical flexure still present. Cephalic flexure less marked. Tail very wide dorso-ventrally. Yolk more elongated, convex on ventral surface, slightly concave on dorsal margin. Pigmentation more pronounced. Mandibular arches coalesced. Boundaries of other arches no longer visible.	Horizontal section shows 44-46 myotomes.	Dorso-ventrally compressed. Yolk granules still present in peripheral portion. Neural arches better developed anteriorly.	Spinal cord shows well defined layer of fibers entirely around it. Layer is thicker in ventral half, thin on dorsal portion. Transverse diameter of cord greater than dorso-ventral. Around central canal a layer of cells radiates from canal. Nerve fibers numerous in ventral roots of spinal nerves.	Pigment increased in posterior layer of optic cup. Retina in three layers. Optic stalk still presents minute lumen; fibers beginning to be formed in its walls. Lens spherical; posterior wall fills entire cavity; epithelium a single layer of cuboidal cells.	Periotic capsule beginning to appear on external ventral side of ear.
32	159 Trans. 160 Sag. 162 Front.	19 mm	43 days		Horizontal section shows 48-50 myotomes.	Anterior end close to hypophysis; slightly knobbed. Portion anterior to heart much smaller than remaining portion; greatly enlarged at level of anterior margin of yolk. Segmentation well marked excepting in caudal portion. Neural arches in anterior trunk region extend dorsally to level of middle of spinal cord.	Three layers well defined in walls of fore brain. Epiphysis a large oval vesicle with thin dorsal wall; communication into third ventricle nearly cut off. Walls of paraphysis irregular. Choroid plexus extending into ventricles. Lateral lobes of infundibulum extended to level of mid brain. Cerebral peduncles formed in ventro-lateral walls of mesencephalon.	Optic cup flattened laterally. Ora serrata better defined. Beginnings of iris and ciliary body. Fibers in optic nerve better developed. Lens cells arranged concentrically; fibers numerous; marked chromatolysis in central portion.	Periotic capsule extends further dorso-mesially.
33	161 Trans. 173 Sag. 174 Front.	20 mm	46 days 2 hrs.	Cephalic flexure reduced. Axis of tail coincides exactly with axis of trunk. Yolk reduced dorso-ventrally, more elongated, more pointed at anterior end. Dorsal and ventral surfaces convex. Pigment better developed.	Horizontal section shows some 50-55 myotomes. In anterior trunk region muscle plate wider, extending ventrally over dorso-lateral margins of yolk. Septa thicker. Two thirds of myoblast fibrillated.	At level of anterior limbs notochord is much larger than spinal cord, at level of posterior limbs much smaller. Yolk not quite absorbed around periphery. Neural arches in anterior trunk region extend dorsally to top of spinal cord, not united; beginning in posterior trunk region; in anterior caudal region indicated in procartilage.	Olfactory nerve connected with fibrous areas in hemispheres.	Layers of retina well defined; pigment more dense. Lumen of optic stalk obliterated near brain wall; exceedingly minute distally; fibers more numerous. In center of lens nuclei have nearly disappeared. Epithelium of lens a layer of cuboidal cells. Fibers well defined, more numerous, concentrically arranged. Eye muscles better developed.	Utriculo-saccular partition extends well into vesicle. Anterior semicircular canal better developed than the others. Macula acustica sacculi well defined. Periotic capsule surrounds ear except mesial surface.

Nose	Hypophysis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gill, Thyroid, Thymus, Trachea, Lung	Urogenital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
Nasal organ connected with surface ectoderm by short stalk. Clear zone in cells much wider. Organ nearly in contact with cerebral hemispheres, separated only by a very thin layer of mesenchyma. Fibrous capsule denser. Tube much longer; narrowed along distal half; stalk extends dorso-caudally; body extends directly caudally. Ventral portion of wall $\frac{1}{4}$ thickness of dorsal.	A solid mass of cells; detached from ectoderm.	Mouth not yet broken through. Cavity well defined but filled with yolk-bearing cells.	Oesophagus occluded. Liver lies for most part on right side of body. Tubules widely separated by sinusoids. Spleen consists of two masses of cells one lying on each side of mesentery.	Gill filaments more prominent. Division of thyroid deeper making two lines of cells at distal portions. Anterior thymus beginning to be detached from gill arches. Distal portion of tracheal evagination lying closely against ventro-mesial portion of anterior end of stomach. Anlagen of right and left lungs in form of tubular evaginations lying in mesenchyma below lateral margins of oesophagus. Tubes extend caudally to level of posterior portion of heart.	Mesonephric tubules coiled. Open into pronephric duct.	Beginnings of trabeculae in ventricle.	Thickened to 1 layer over head. Glands and sense cells developed. Few pigment cells in epidermis.	Trabeculae extend forward to optic nerve; curved downward at anterior end; caudally, are continuous with basal plate; at level of posterior margin of ear they come together at sides of notochord and grow up on each side of medulla. Labial cartilages small, united in median line. Anlage of second basibranchial. Earliest traces of ossification in base of skull. Layer of condensed mesenchyma around nasal pits. Otic capsule extending medially over dorsal and ventral surface of ear.	Anterior limbs short stumps.	30
First indications of olfactory nerve.			Evaginations in wall of stomach indicate beginning of glands. Gall bladder shows as well defined vesicle with columnar epithelial walls.	Gill bars much elongated. Gill filaments better developed. Lungs show now as pair of lateral tubular diverticula; epithelium low columnar, nuclei occupying peripheral position. Anlage of pleura as layer of cuboidal cells. Some mesenchyma between lungs and oesophagus.	Distal ends mesonephric tubules have walls thinner and infolded. Ureters very small.		Cells of corium slightly flattened.	Walls of otic capsule chondrified except on median and dorsal sides. Neurapophyses appearing in anterior portion of trunk region. Pectoral girdle chondrified; narrow scapular portion extending to level of notochord; larger ventral portion (coracoid) about half as long as scapular. Humerus chondrified; radius and ulna in procartilage.	Anterior limbs 1 mm long; portion of trunk project dorsocaudally. Posterior limbs .5 mm long.	31
		Mouth a deep transverse groove. Cavity still filled with cells.	Lumen of stomach sharply marked. Epithelium long columnar cells, still laden with yolk granules. Muscular walls beginning to be formed. Yolk nearly absorbed in posterior end of cloacal walls. Liver further developed; contains larger number of tubules more compactly arranged. Sinusoids smaller. Gall bladder larger and better defined. Cystic duct and ductus choledochus formed. Dorsal pancreas narrower, its lumen, where it enters gut, constricted to form a duct. Anlagen of ventral pancreases appear as dorso-lateral evaginations of ductus choledochus.	Thyroid divided into two portions which are seen in transverse sections as small groups of cells lying just dorso-laterally from the sterno-hyoid muscle. Lungs slightly longer.	Pronephros beginning to degenerate. Mesonephros extends over 8 segments, beginning with the 9 <sup>th</sup> . Not segmentally arranged. 15 tubules.		Earliest skin glands in mid dorsal region. Sense organs developed.	Beginning differentiation of hypohyal and ceratohyal, also of first hypobranchial and ceratobranchial. Second basibranchial developed. Antorbital process formed. Condensed mesenchyma indicates position of inner nasal plate. Anterior ends of MECKEL's cartilage united by procartilage; also lateral margins of posterior arches. Occipital processes fused with wall of otic capsule.		32
Organ further elongated caudally. Depressions forming around exterior indicating beginning of lobulation. Capsule not so well defined as in preceding stage. Organ connected with anterior end of foregut by a solid cord of cells. Pigment beginning to grow around stalk. Olfactory nerve in connection with brain.	Greatly elongated transversely.	Groove deeper. Anlage of teeth. Anlage of tongue.	Boundary between stomach and intestine sharply marked owing to greater absorption of yolk in epithelium of stomach and formation of muscular wall of stomach. Intestine now much coiled. Cloaca differentiated from intestine by greater absorption of yolk. Oesophagus occluded. Faint sacculations indicate beginning of glands in wall of stomach. Hepatic cords widely scattered, separated by large sinusoids. Cells filled with yolk granules. Gall bladder small, lined with columnar epithelium. Empties by cystic duct into ductus choledochus. Dorsal pancreas well differentiated; duct larger except where it enters intestine. Ventral pancreases contain few tubules. Their two short ducts join cystic duct as it emerges from liver to enter ductus choledochus.	Gills longer; many filaments. Thymus, first and second portions detached from gill arches, third still connected. Lungs longer, further evaginated, lumina larger. Epithelium low columnar, heavily laden with yolk granules. Tips reach nearly to level of anterior margin of liver.	Pro-nephric tubules still open into coelom. Anlage of bladder as slight evagination of ventral wall of cloaca.	Auricles beginning to show bilobed condition. Walls of ventricle very thin; very few trabeculae.	Skin glands in dorsal region well developed. Corium cells flattened. Basement membrane formed.	No cartilage in pelvic girdle. Anlage of neurapophysis throughout trunk region, none in tail. Neural arches in anterior portion extend dorsally to top of spinal cord; not united. Caudally neural arches indicated in procartilage. Ossification beginning around extremities of MECKEL's cartilage.	Anterior limbs project caudally; Anlage of digits. Posterior limb buds pointing caudally.	33

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
34	175 Trans. 180 Sag. 181 Front.	21 mm	49 days	Axis of head, trunk and tail coincide with line slightly arched upward over yolk. Yolk elongated, narrowed dorso-ventrally; narrowest portion anterior; viewed from above pear-shaped, narrow end anterior. Tail very wide dorso-ventrally. Irregular band of pigment along side of body. Anterior limbs project postero-ventrally three digits formed. Posterior limbs project caudad.		Caudal portion filamentous with slight knob at extreme end; still rests against neural tube. Mesenchymal thickenings in sheath present in caudal portion.	Flexures less marked. Hemispheres extend far in front of dienecephalon. Epiphysis and paraphysis large. Plica rhombomesencephalica well defined. Lamina cerebellaris short; commissura superior and posterior and habenularis all well developed. Spinal ganglia very large. Nerves well developed.	Ganglionic layers of retina marked off from inner layer; layer of rods and cones indicated by layer of elongated cells. Pigment prominent in iris. Epithelium of lens flat on outer surface, cuboidal on inner.	
35	182 Trans. 183 Sag. 184 Front.	22 mm	52 days			Anterior end of notochord separated from hypophysis by thin layer of mesenchyma. Surrounded by very dense membrane. Slightly constricted at level of mesenchymal condensations indicating boundaries of vertebrae. Indications of ossification in sheath. Anlage of haemal arches in anterior portion of tail.	Paraphysis shows few lateral diverticula. Ganglion habenulare and tuberculum posterius well defined. Commissura cerebellaris formed, also commissura anterior and commissura pallii anterior. Layer of fibrous matter around cord thicker. Band of cells on dorsal border of cord larger, dorsal half of central canal nearly closed. Pia mater better defined.	Rods and cones recognizable. Iris more deeply pigmented; lies close against epithelium of lens. Optic nerve well differentiated; still contains minute lumen near eye. Anlage of cornea as single layer of mesenchymal cells.	Utriculo-saccular partition extends about half way across vesicle. Semicircular canals partially enclosed in cartilage. Periotic capsule entirely surrounds ear.
36	186 Trans. 188 Sag. 191 Front.	23 mm	55 days			Does not quite reach posterior end of neural tube. Yolk absorbed except at extreme anterior end. Sheath strongly developed. Constrictions better marked. Neural arches developed throughout body; meet dorsally in trunk region but not in tail. Ossification beginning in anterior neural arches. Haemal arches better defined in tail. Where neural processes are given off notochord is compressed laterally giving it a triangular outline in cross section.		Rods and cones better defined; outer and inner nuclear layers indicated; ganglionic layer well defined. Pigment dense in retina and iris; some pigment in choroid. Lens epithelium flatter. Sclerotic indicated in band of dense mesenchyma.	

Nose	Hypo-physis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
Organ longer, becoming slightly concave on ventral surface. Further indications of lobulations.		Mouth cavity still closed.	Glands better defined. No differentiation into pyloric and cardiac glands. Liver roughly triangular in section, has extended caudally. Gall bladder still small, lining still a single layer of columnar cells filled with yolk granules. Hepatic cords more definite owing to absorption of yolk. Dorsal pancreas lumen and duct smaller. Several small tubular evaginations coming off from it. Ventral pancreas lobules not distinct. Spleen an elongated mass of cells compressed dorso-ventrally lying in mesentery just above stomach.	Gills further developed showing greater number of filaments. Gill clefts open between first and second arches and between second and third branchial arches but fifth cleft not open. Lumina of lungs still larger. Tips of lungs reach anterior margin of liver.			Cells of stratum germinativum flattened. Glands and sense organs more numerous.	Procartilage present in pelvic girdle and in femur. Ossification in side walls of brain case, also in base. Anlage dentale as ossified plate on external surface of MECKEL'S cartilage. Beginning of angulare thin plate of bone. Indications of mento-mandibulare. Ossification beginning in premaxilla and in quadrate.	Anterior limbs project ventrally; digits plainly defined. Posterior limb buds project caudad.	31
Nasal duct opens into pharynx on one side. Connective tissue and blood vessels extend into ventral wall giving it a much lobulated appearance; dorsal wall also slightly lobulated.		Mouth broken through.	Dorsal pancreas smaller, tubules fewer. Yolk absorbed in cells. Prozymogen granules abundant.	Trachea larger and longer. Lungs longer, caudal ends behind heart, over anterior portion of liver. Walls better defined, epithelium cuboidal. Yolk nearly all absorbed.	Glomeruli well defined. Tubules very much coiled.	Auricles further developed. Cavity of ventricle large, walls thin, many trabeculae. Walls of truncus much thicker than those of conus. Valves beginning to be formed between conus and ventricle; also between conus and truncus.		Occipital arch growing up from sides of notochord; below uniting posterior ends of parachordals. Parachordals fused with basal plates. Trace of trabecular crest to which process of quadrate is attached. Short otic process fused with otic capsule. Coracoid process extends nearly to ventral margin of liver; scapular extends to upper margin of notochord. Humerus larger; glenoid cavity very deep. Ulna and radius chondrified. Anlage of carpals and metacarpals. Pelvic girdle chondrified; consists of dorsal iliac portion and ventral pubic portion. Femur chondrified; anlagen of tibia and fibula.		35
Nasal tubes open into pharynx on both sides.		Posterior median portion of cavity still contains yolk-bearing cells. Papillae of teeth formed.	Posterior portion of liver has extended dorsally to level of posterior portion of stomach; contains few bile ducts. Cells loaded with yolk. Tubules much more compact; sinusoids narrower, changing to capillaries. Gall bladder larger, cells in wall low columnar or cuboidal, free from yolk. Dorsal pancreas 6-8 tubules. Ventral as above. Spleen larger, more vascular.	Trachea smaller, lining layer of cells flat. Lungs much farther distended laterally, extending to extreme lateral margins of the liver. Pleura consist of layer of closely apposed cuboidal cells.	Pro-nephric tubules reduced in size.		Glands larger; sunk below level of epidermis; contain lumen. Cells of stratum germinativum elongated flattened.	Floor of otic capsule nearly complete. Ossification beginning in parasphenoid also in parietals and frontals. Anlage of vomer. Neural arches developed throughout body; meet dorsally in trunk, not in tail. Haem-apophyses appearing in anterior portion of tail. Ossification beginning in neural arches. Bone well developed around centers of vertebrae. Traces of ossification in scapula and humerus.		36

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
37	192 Trans. 195 Sag. 167 Front.	24 mm	58 days			Anterior end in contact with parasphenoid but does not reach level of hypophysis. Constrictions opposite bases of neural arches more marked anteriorly. Ossification well marked around sheath. Haemal arches approximating ventrally in tail. Anlage of lateral basal process in cartilage in anterior portion of body.	Paraphysis with many lateral diverticula; still opens into third ventricle. Eminentia Pallii medialis striatum (?). Pallium and subpallium differentiated. Septum ependymale present. Beginning of taenia fornicis.		
38	198 Trans. 199 Sag. 200 Front.	25 mm	61 days	Flexures as above, yolk elongated, diameter reduced dorso-ventrally and laterally. Tail broader, less pointed. Surface more deeply pigmented; bands more marked. Both anterior and posterior limbs possess four digits. Pigment bands more sharply defined.	In anterior trunk region wider occupying four fifths of space between notochord and skin; extended ventrally to lower margin of yolk. Myoblasts in upper portion of myotome completely filled with fibrillae.	In anterior portion much compressed laterally, more compressed at bases of neural processes. At level of anterior limbs smaller than spinal cord; in trunk region and at level of posterior limbs large as cord; in tail about twice the diameter of cord. Yolk all absorbed. Haemal arches well developed, free ends coalesced. Ossification very considerable in neural arches and notochordal sheath. Lateral basal processes better developed throughout trunk region.			Ductus endolymphaticus extends over dorso-lateral walls of medulla; stalk very narrow but still widely open into sacculus. Semi-circular canals better developed; all partially enclosed in cartilage. Lagena a more extended evagination. Anlagen of pars neglecta and macula acustica neglecta. Macula acustica sacculi two layers of cells thick.
39	201 Trans. 204 Sag. 205 Front.	26 mm	64 days 4 hrs.			Compressed dorso-ventrally in anterior part of body, laterally in posterior part. Anterior end surrounded by bone.	All divisions of brain clearly defined. Plexuses extending from hemispheres to medulla. Optic nerves very small, hollow near brain. Mid brain short, slightly arched. Ganglia habenulae large, symmetrical. Velum transversum contains numerous large blood vessels. Infundibulum wider than mid brain. Cerebellum extremely small. Spinal cord oval in section. White matter on outside thickest in lateral portions. Dorsal columns large, broader anteriorly. Central canal very small nearly circular in section. Some granules still present in cells of cord. Pia mater well developed.	Eye not so much flattened laterally. Retina shows layer of rods and cones, outer nuclear layer, outer reticular layer, inner reticular layer, ganglionic layer. External limiting membrane well defined. Choroid better differentiated; pigment layer closely apposed to pigmented layer of retina. Lens spherical, surrounded by a capsule of a single layer of flattened cells. Fibers well developed. No nuclei in center. Posterior chamber contains a few mesenchymal cells. Corneal epithelium in two layers. Eye muscles well developed.	

Nose	Hypo-physis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
Wall of nasal duct consists of two layers of cuboidal cells.	Lies against floor of skull surrounded by a connective tissue capsule. More lobulated. Blood vessels growing into it.	Entire mouth open. Oesophagus open. Teeth projecting slightly into mouth cavity, ossified. Tongue better developed.	Glands in stomach well formed. In anterior half numerous, flask-shaped; in posterior half tubular. Both kinds unbranched. Yolk absorbed in epithelial cells of stomach also in hindgut; midgut cells still heavily laden. Oesophagus open. Liver extends further caudally and dorsally. Hepatic cords more closely approximated. Yolk partially absorbed. Both dorsal and ventral pancreases larger, showing greater number of tubules. Spleen lies pressed against dorsal wall of body cavity; surrounded by thin layer of connective tissue; shows numerous dividing cells; much connective tissue.	Trachea longer reaching to caudal region of heart before dividing. Lungs longer, walls better defined, still very thick. Caudal ends detached from intestinal wall except for short thin mesentery. Epithelium becoming flattened. Mesenchyma less compact. Lungs extend over anterior two thirds of liver laterally to dorso-lateral margins of liver. Posteriorly attached by band of mesenchyma to lateral margins of wall of stomach.	Pronephros reduced in size. Tubules cut off from coelom? Duct (ureter) opens into cloaca. Mesonephric tubules greatly convoluted; extends from 9" to 17" somite. Bladder comes off from cloaca near point where ureters enter. Sexual cells, a single row of cells containing large yolk granules. Opposite 15" segment.		Glands appearing in ventral region, anteriorly and posteriorly; numerous in lateral portion. Sense organs more numerous.	Supra-scapula well formed in cartilage extending dorsally to level of spinal cord. Tibia and fibula chondrified; no tarsals or metatarsals. Ilium and pubis larger. Mesenchyme around nasal pits. Ossification beginning around margin of trabeculae. Parasphenoids forming as long plates on lower margin of basal plates extending forward to region of eye. Vomer ossified. Frontal ossified at sides and posterior end. Neural spines beginning to appear anteriorly. Indications of ossification in femur. Ossification beginning in neural processes and in haemal arches.		37
			Yolk nearly absorbed in liver.	Gills more prominent; filaments much branched. All clefts perforate excepting fifth. Lateral portion of thyroid contains small irregular vessels. Pleura proximally layer of flattened cells. Some mesenchyma between pleura and epithelium of lungs.			Oval cells and sense organs more numerous. Sensory hairs evident. Stratum corneum very flat cells. Stratum germinativum two layers.	Basilar plate continuous with capsular floor and trabeculae. Jugular foramen formed, elongated. Otic process of quadrate united with otic capsule. Antorbital process further developed. Haemapophyses well developed; not united ventrally. Ossification in phalanges of anterior limb, also in iliac process.	Limbs better defined; anterior reach ventrally to border of yolk. Digits better defined in both.	38
Antero-lateral wall very thin. Lobulations prominent. Epithelium of dorsal wall ciliated. Olfactory nerve clearly defined.	Enlarged transversely. Divided into two portions composed of cells of different staining capacity, anterior part more deeply stained. Posterior part more lobulated and more vascular.	Unicellular glands in epithelium of mouth and tongue. Teeth extending into mouth cavity. Tongue well defined.	Unicellular glands present in mouth, pharynx, oesophagus and hindgut. Extreme anterior portion of stomach contains no glands. Pyloric glands much branched. Intestine lies in 4 to 5 transverse coils. Cavity of small intestine widely open. Muscular walls better developed. Epithelium consists of single layer of long columnar yolk-laden cells. Hindgut larger; yolk absorbed; anterior part shows evaginations (glands of LIEBERKÜHN?). Liver lies closely in contact with dorsal wall of coelomic cavity. Yolk all absorbed. Gall bladder very large, surrounded by a layer of connective tissue, epithelium flat. Dorsal pancreas has grown caudally and contains 50 or more tubules; ventral contains about 12 tubules, is about $\frac{1}{3}$ size of dorsal. Ducts of both lined with cubical epithelium.	Thymus consists of three spherical epithelial patches on each side. Two lie close together opposite posterior portion of ear. Anterior largest; posterior much the smallest. Trachea much larger, walls thinner, epithelium flatter. Lungs much larger, tapering caudally to end at level of posterior limbs; are attached to intestinal wall and to liver by short mesentery. Left lung larger than right and extends further caudad.	Pronephros degenerating. Extends from middle of 3" segment to middle of 4", hanging down into body cavity. Duct very small. No openings into body cavity discernible. Mesonephros extends from 15" to 20" segments. Ducts open into cloaca. Bladder much folded, opening into cloaca by short urethra, surrounded by connective tissue. Sexual glands two rods of large oval cells extending from 13" to 18" segment, just ventrad of mesonephros.	Auricles nearly same size in transverse section. Sinus venosus very large. Truncus arteriosus divided longitudinally. Trabeculae appearing in walls of auricles. Marked constriction between sinus and auricles.	Skin glands present in abdominal region. Stratum corneum a layer of squamous cells. Basement membrane dense.	Internasal cartilage connects trabeculae anteriorly. Pubic bones united in median line. Tarsals and metatarsals chondrified. Palatopterygoid and sphenoid ossified in posterior portion.		39

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
40	206 Trans. 207 Sag. 209 Front.	27 mm	67 days			Chorda more compressed laterally at points where neural arches come in contact. In trunk region and in region of posterior limbs transverse diameter equals that of spinal cord; in tail region diameter twice as great as that of cord. Beginning ossification in haemal arches.			Ear much extended antero-posteriorly compressed dorso-ventrally. Ossification beginning in external portion of otic capsule.
41	210 Trans. 211 Sag. 212 Front.	28 mm	70 days 4 hrs.	General contour of body much changed, owing to rapid absorption of yolk. Tail broader dorso-ventrally. Constricted at level of posterior limbs. Yolk very long oval, flatter on ventral surface. Pigmentation extending down over one half of lateral surface of yolk.					
42	213 Trans. 214 Sag. 216 Front.	29 mm	74 days 12 hrs.						
43	217 Trans. 218 Sag. 219 Front.	30 mm	78 days		In anterior trunk region myotomes wider occupying nine tenths of space between notochord and skin; extend ventrally to median line where they coalesce.	Much constricted in region where cartilages are in contact with body of vertebrae. Thickenings of sheath to form intervertebral disks better marked. Cartilages lie between layer of bone and sheath. Ossification beginning in lateral basal processes. Cartilaginous ribs present.	Brain and cord nearly straight line. Groove present between diencephalon and mesencephalon. Deep groove in front of cerebellum. Plexuses large filling greater portion of 3 <sup>rd</sup> ventricle extending into lateral ventricles and into 4 <sup>th</sup> ventricle. Distal end of paraphysis lies on level with epiphysis. All commissures better defined. Brain fills brain case at anterior end, not posteriorly. Spinal cord does not nearly fill canal. Dorsal columns of cord well defined; gray commissure present between them and gray matter of cord. Central canal very small, circular in section.		Endolymphatic sac wider; duct narrower. Cristae acusticae of canals formed. Macula acustica sacculi a thickened area of epithelium three rows deep. Semicircular canals enclosed in cartilage. Ossification more extended on lateral surface of periotic capsule.

Nose	Hypophysis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
			Intestine crosses abdominal cavity 6-7 times. Dorsal portion of liver connected by narrow band with ventral portion. Hepatic cords closer together. Few hepatic ducts. Dorsal and ventral pancreases not in contact yet. Tubules more numerous in dorsal.	Lungs reach to level of caudal end of liver.			Unicellular gland and sense organs very numerous. Sensory hairs projecting beyond surface.	Ossification beginning in haemal arches.		40
				Walls of trachea and lungs very thin; surrounded by mass of mesenchyma. Lungs flattened against stomach. Epithelium thinner; pleural covering very thin.				Ossification well defined in haemal arches.		41
			Caudal portion of liver occupies over one half of coelomic cavity. Numerous hepatic ducts. Dorsal and ventral pancreases in contact. Dorsal much the larger. Spleen longer; compressed dorso-ventrally between intestine and dorsal body wall. Connective tissue more abundant.	Gills better developed, pigmented. Filaments greatly increased in number. Second, third and fourth gill clefts widely open to exterior; first and fifth imperforate.					Limbs longer, pigmented, both pairs functional. Four digits well defined on each.	42
		Under lip marked by a very deep transverse groove. No upper lip. Tongue being cut off on ventral surface by deep lateral constriction.	Extreme anterior end of stomach contains no glands except unicellular. In anterior portion flask-shaped glands are larger and more numerous. Middle and posterior portion contains branched tubular glands. Intestinal epithelium better defined; still contains yolk granules. Hindgut contains branched glands. Long process of liver extending posterior to gall bladder; another in median line extending over dorsal wall of gut.	Right lung lies against margin of liver; caudal end in contact with spleen.	Pronephric ducts small but still open. Mesonephros begins 3 or 4 segments behind pronephros and extends over about 10 segments. Bladder epithelium short columnar cells, enclosed in connective tissue. Walls much folded. Sexual glands extend from 14" to 20" segments. Yolk granules present in posterior portion.	Pericardium well defined, very thin, detached from muscular wall. Auricles widely in communication with each other. Blood passes into ventricle by single opening on left side. Valves present in this opening. Semilunar valves between ventricle and conus appear as extended proliferations of wall. Valves also present between conus and truncus.	Mucous glands larger, abundant, none on ventral surface. Unicellular glands more numerous.	All the cartilages of carpus and tarsus metacarpus and metatarsus and phalanges formed. Middle portion of MECKEL'S cartilage surrounded by bone. Pre-maxillae and frontals well ossified. Cartilage appearing in center of body of vertebrae.	43	

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
44	220 Trans. 221 Sag. 223 Front.	31 mm	82 days 12 hrs.						
45	224 Trans. 225 Sag. 226 Front.	32 mm	87 days						
46	227 Trans. 229 Sag. 230 Front.	33 mm	92 days						
47	231 Trans. 232 Sag.	34 mm	97 days	General form of body resembles adult. Ventral surface of head and trunk nearly in straight line. Anterior and posterior limbs project far below ventral surface of body. Pigmentation bands more precisely marked. Dorsal median dark band, below this a light band, then a broad dark band.					

In caudal region larger than spinal cord. More constricted in centers of vertebrae. Intervertebral cartilages more prominent. Layer of bone thicker.

Bone surrounding anterior end thicker. Ribs better developed; extend one half of distance to lateral surface of body. Neural arches at base more completely ossified.

Nose	Hypophysis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Skeleton	Limbs	Stage
			Liver still larger in caudal portion occupying $\frac{3}{4}$ of body cavity. Has several hepatic ducts entering cystic duct. Gall bladder removed some distance from ventral pancreas. Dorsal and ventral pancreases have united into a continuous mass.	Tip of lung at caudal end of liver. Dorsal walls of lungs in region anterior to liver very much thinner than ventral wall.						41
Compressed anteriorly between infundibulum and floor of skull; broader posteriorly. Much lobulated.			Lumen of intestine larger; epithelium a single layer of poorly defined cells. Muscular wall very thin, a single layer of cells. Few hepatic ducts seen in body of liver; a few enter the cystic duct. Dorsal pancreas tubules compactly arranged.	Lungs very much larger, owing to thinness of walls very much collapsed. Tips of lungs still show embryonic condition. Cartilages formed in walls of trachea. Epithelium of lateral walls of trachea much thicker than dorsal and ventral walls. Trachea in anterior portion shows columnar cells; ciliated? At level of heart epithelium flat.	Pronephros compressed dorso-ventrally; extends over one segment. Mesonephros extends over about 10 segments.		Unicellular glands extremely numerous.			45
			Posterior portion of liver divided by transverse fissure into smaller dorsal and larger ventral portion. Gall bladder very large; extends far caudal of liver; lined with flattened epithelium. Cystic duct larger and longer. Pancreas large extending around dorsal and lateral portion of intestine. Ducts open into cystic duct; show numerous branches with tubules emptying into them. Spleen larger, more vascular.		Sexual glands extend far forward; begin to project down into body cavity; more distinctly separated from mesonephros.					46
			Intestine forms 6 or 7 transverse folds. Intestinal cavity very large. Muscular wall of midgut consists of two or more layers of cells. In hindgut unicellular glands very numerous. Dorsal and ventral pancreases more intimately united; dorsal larger and contains greater number of tubules. Spleen lies just to left of median line in concavity in dorsal wall of stomach.	Tracheal cartilages better developed at anterior end. Lungs on either side of stomach. Walls irregularly folded. Left lung still notably shorter than right and lying nearer mesial plane of body.	Pronephros smaller; lumina of many tubules obliterated; duct degenerating. Mesonephros extends from 9" or 10" segment over about 13 segments. Müllerian duct not formed yet.		Scapula flattened laterally, broad antero-posteriorly. Coracoids meet in mid-ventral line. Ossification beginning in pectoral and pelvic girdles, also in humerus and femur. Sphenoid ossified. Parietals ossified in median portion. Opisthoticum ossifying, also occipital arch.	Limbs much longer. Digits longer.		47

Stage	Series	Length	Age	Body Form	Somites	Notochord	Nervous System	Eye	Ear
48	233 Sag.	36 mm	110 days			Anterior end at level of hypophysis, sometimes beyond it. Intervertebral cartilages thicker. Bodies of vertebrae more constricted. Layer of bone around cord thicker.			
48a		38 mm							
49	234 Trans. 236 Sag. 237 Front.	39 mm	126 days	Has practically reached adult condition except in coloration. Same bands prominent as in preceding stage. Lateral band mottled.		Ossification very complete in neural spines and basal processes in ribs, and in haemal arches; beginning in haemal spines.		Eye nearly spherical, slightly flattened laterally. Retina as in 26 mm excepting ganglionic layer is one cell thick. Other layers together with external and internal limiting membrane sharply defined. Iris, choroid, cornea and sclerotic as in 26 mm. Lens capsule cells much more elongated tangentially.	

Nose	Hypo-physis	Mouth	Digestive System, Liver, Pancreas, Spleen	Gills, Thyroid, Thymus, Trachea, Lung	Urino-genital System	Heart and Blood Vessels	Skin	Skeleton	Limbs Stage
			<p>Epithelium of pharynx consists of 2 or 3 layers of flat cells containing numerous unicellular glands. Oesophagus thrown into numerous longitudinal folds; epithelium columnar, ciliated, with unicellular glands. Muscular wall very thin. Liver very large. Lies on right side. In frontal section outline is long triangle, in transverse section an equilateral triangle. Posterior end shows several irregular processes. Pancreas elongated cephalo-caudally. Anterior portion flattened and closely applied to dorso-mesial wall of intestine. Posteriorly greatly thickened and roughly triangular in shape. Posterior end lies on right side.</p>						48
			<p>Anterior wall of stomach thin; circular band of muscular fibers very thin, increasing in thickness toward pylorus. Epithelium columnar. Submucosa developed. Extreme anterior end free from glands. Flask-shaped glands confined to anterior portion. Enlarged part of glands consists of a single layer of very flat cells; duct, cuboidal or columnar. Middle and posterior portion of stomach contains branched tubular glands. In pyloric portion epithelium is thrown into longitudinal folds. Inner ends of cells form a striated border which extends into intestine. Liver contains numerous bile ducts. Hepatic cords well defined. Highly vascular.</p>						48 a
Compressed; extended laterally. Contains many tubules.	Teeth project far into mouth cavity. Tongue narrower in front. Taste buds well developed in epithelium of tongue.	<p>Intestine shows strong circular and longitudinal bands. Walls thrown into longitudinal folds. Lumen greatly expanded at level of posterior limbs forming rectum? Epithelium columnar. Unicellular glands more numerous posteriorly. Gall bladder large oval, extends caudally beyond liver. Walls formed of a single layer of flat cells. Cystic duct large. Pancreas composed of branched tubules. Dorsal portion opens by a single duct just caudally of stomach; ventral portion smaller, opens by two ducts. Spleen lies on left side, in frontal or sagittal section long, oval pointed in front. Arrangement of cells suggesting tubules.</p>	<p>Gills all developed. Second, third and fourth arches perforate, first and fifth not broken through. Thyroid small, halves widely separated; consists of 20-30 large vesicles. Lies close against first epibranchial. Many vesicles filled with colloidal material. Trachea at caudal end much larger than oesophagus. Divides at level of glenoid cavity. Lungs much longer extending some distance behind liver. Right lung larger than left and extends further caudad. Epithelium very thin and flat.</p>	<p>Pronephros degenerating. Lumina of many tubules obliterated. Duct degenerating. Mesonephros extends from 9<sup>th</sup> or 10<sup>th</sup> segment over about 13 segments. Müllerian duct not formed. Bladder large, wide transversely, thin-walled, extending well forward beyond pelvis. Sexual glands nearly coextensive with mesonephros.</p>	<p>Heart well developed. Auricles well defined folded over ventricle. Imperfect ventricular septum. Ventricle with numerous muscular trabeculae. Truncus arteriosus thick walled; conus thin. Branchial arteries well developed. Pulmonary arteries and veins very small. Hepatic veins very large; they unite to form sinus venosus which empties into the right auricle. Internal and external jugulars present; jugular sinus well marked. Splenic, gastric and mesenteric veins formed. Caudal vein gives off branches to form renal portal system. Arteries and veins in limbs complete.</p>	<p>Internasal cartilage well defined. Lateral and dorsal wall of brain capsule now a thin layer of bone. Bone in whole extent of lower jaw. Nasal capsule formed in cartilage. Hypobryal much larger. Second hypobranchial still very small. Cartilage appearing around orbit. Membrane bones of skull generally formed. Tibia and fibula, tarsals and metatarsals ossified.</p>	49		

### **Some Variations in External Structures.**

Before the individual variations are considered in detail it should be stated that a given nest contains but few eggs that are in precisely the same stage of development. The differences are most obvious in the early stages up to the closure of the neural fold. From the closure of the neural folds to the 39 mm larva the variations are not so pronounced, yet there are innumerable minor variations.

In the following descriptions the principal variations observed in the external features are first recorded, then those observed in the sections.

#### **Cleavage.**

In the cleavage stages, from the second on to late cleavage, there is so much variation in the position, extent and rate of progress of the various grooves that it is impossible to record them. Some of the variations have been described elsewhere (1904 b) by the senior author.

#### **Gastrulation.**

Some variation is found in the position of the first line of invagination which forms the dorsal lip of the blastopore. Its first appearance may be along a line equidistant from the equator and the vegetative pole or it may form nearer the equator and again sometimes nearer the vegetative pole. The first line of invagination may be nearly straight and again it may be decidedly crescentic. The maximal diameter of the yolk plug may equal one half the diameter of the egg. The closure of the blastopore usually occurs in about six days but it may close in five days.

#### **Closure of neural folds.**

Variations in the closure of the neural fold are frequently observed. The coalescence of the lateral portions of the fold usually begins at the posterior end. Sometimes they first coalesce along the middle portion of the embryo. Usually the transverse portion of the fold is continuous; at other times it shows a deep transverse groove which separates it into right and left halves. Cf. Figs. 18, 19, 20, 21. Wide variations exist in depth, width and extent of the neural groove.

#### **Appearance of optic vesicles.**

The optic vesicles are usually present before the complete closure of the neural fold. Sometimes they are present as disc-like depressions in the neural plate before the lateral portions of the fold have begun to coalesce; again, but rarely, they are not visible until the folds have closed.

#### **Neuromeres (?).**

There are frequently observed in the cephalic region well marked serial elevations and depressions along the inner margins of the lateral folds and across the neural plate. In other embryos there is not the slightest trace of either.

### Somites.

In some embryos three myotomes are differentiated before the neural folds are closed, while in others there are no traces of myotomes until the folds are closed. The number of myotomes early becomes exceedingly variable in the tail. It is here impossible to count them accurately either in surface views or in sections. In the trunk i. e. between the limbs from the 15–16 mm larva up to the 39 mm they seem to be fairly constant numbering 18–20. In the tail however they are variable, so that in larvae of identical lengths there may be a variation of 1–5 myotomes. It should be emphasized that the number in the tail is determined with great difficulty since the most caudal are but slight thickenings in the mesoderm.

### Lateral curvature of body.

In the embryos of 8–15 mm there is much variation in the lateral curvature of the body. In some nests as high as 80% of the embryos have the head and tail curved to the right. Other nests show a like percentage in which the head and tail are curved to the left. Out of 328 eggs, taken from five nests, 174 had the concave side on the right and 154 on the left.

### External gills.

In most embryos of 9 mm (Fig. 25) the fourth and fifth arches are a common mass with no indications of the line of division; in some the line of invagination is distinct. In the later stages (e. g. 25 mm) there are usually five or six filaments on the middle gill bar, in others there are eight or nine. This variation is even more pronounced in the 26–39 mm larvae.

### Limbs.

Some variations have been noted in the time of appearance of both the anterior and the posterior limb buds. The anlage of the anterior limb is usually discernible in the 11 mm stage (Fig. 27), but sometimes not until the larva measures 12 mm. The posterior limb buds are usually beginning in the 12 mm stage, sometimes are not present until the larva measures 13 mm. In the formation of the digits variations are found. Three are usually present on the anterior limbs in the 20–21 mm larva. In some three are not present until the larva is 23 mm long. The same variation is observed in the time of formation of the fourth digit on the anterior limb. Similar variations are found in the time of formation of the posterior digits.

### Pigmentation.

Although little variation is observed in the position of the bands there is much variation in the degree of pigmentation. These differences are most pronounced in the larvae between 19–25 mm. In some the chromatophores are densely aggregated while in others they are sparsely scattered. In some (25 mm) they have extended over the dorsal portion of the yolk only, while in others they have extended over one half of the lateral surface of the yolk.



### Variations in Internal Structures.

In the study of the variations of internal structures only those which are most obvious have been recorded.

#### **Notochord.**

There is considerable variation in the anterior extent of the notochord in nearly all stages. This variation is more obvious in the earlier than in the later stages. Usually a considerable tract of mesenchyma lies between the anterior end of the chorda and the hypophysis. Frequently this tract is short and sometimes only a narrow band.

#### **Eye.**

In sections the optic vesicles show the same variations which have been recorded under variations in external features. The appearance of pigment in the retina and iris shows considerable variation. In the 17 mm embryo series 140, 141, 145 show considerable pigment in the retina and its first appearance in the iris, while 146 shows the beginning of pigment in the retina and none in the iris. The time of first appearance of pigment in the choroid is likewise variable. In the 18 mm embryo series 152, 155 show considerable pigment, series 146, 151 show but little pigment, while in series 147, 149, 150 there is no pigment in the choroid.

#### **Ear.**

In the 6 mm embryo, series 100, there is a cup-shaped invagination of the inner layer of the ectoderm while in series 101 there is a vesicle. In the 7 mm embryo, series 102, shows the otic vesicle just closed, while in series 103 it is not quite closed. In the 18 mm larva, series 146 and 147, the endolymphatic duct extends over the medulla to about one half the distance to median line; in series 151 it extends over the lateral margin of the medulla; in series 149 it does not extend over the lateral margin of the medulla.

#### **Nose.**

In the 9 mm stage series 100 shows the anlage of the nose as a shallow invagination of the ectoderm. In series 107 there is no invagination. In 108 it is slightly cup-shaped. In series 109 thickened ectoderm, in series 110 thickened ectoderm.

#### **Epiphysis.**

In the 9 mm stage series 106, 109, 108 show the epiphysis to be a slight cup-shaped evagination in the dorsal wall of the fore-brain. In series 107, 110 the evagination is deeper and somewhat pear-shaped.

#### **Paraphysis.**

In the 11 mm embryos, series 116, the paraphysis is a very shallow evagination in the postero-dorsal wall of the telencephalon. In 117 the paraphysis is a deep pit-like evagination.

### **Hypophysis.**

In the 9 mm embryo, series 106, 107, the anlage of the hypophysis is two layers of cells thick, in 108 it is but a single layer, while in 109 it is three or four layers thick.

### **Liver.**

In the 9 mm embryo, series 106, 107, 108, 110 the anlage of the liver is shown as a wide evagination of the gut; while 109 shows two or three tubules.

### **Pancreas.**

The dorsal pancreas seems to grow very slowly and to show much variation. Series 118 and 119 show the dorsal pancreas as a slight evagination of the dorsal wall of the gut. In series 120 it is in about the same stage. In series 125 the evagination is vesicular. In the 13 mm embryo series 121, there is a cup-shaped evagination. In 122, 124 the evagination is vesicular. In the 14 mm stage it varies from cup-shaped evagination in series 127, 130 to vesicular in 129.

### **Spleen.**

In the 15 mm embryo, series 131, 134 show the anlage of the spleen as a small mass of mesenchymal cells in the dorsal mesentery of stomach. In series 132 there is no indication of spleen. In 133 there is a well defined mass of mesenchymal cells.

### **Thymus.**

In the 16 mm embryo, series 137, 138, 139 the thymus begins as a small mass of cells in close proximity to the 3rd branchial cleft. In 136 the thymus anlage is not yet present.

### **Trachea—Lungs.**

In the 10 mm embryo series 111, 112 show the beginning of the trachea as a short groove in the ventral wall of the pharynx. Series 113 shows a deeper vesicular evagination.

### Bibliography.

The literature comprises the papers on those forms which belong to the Ichthyoidea (CLAUS) including the Perennibranchiata (*Proteus*, *Siren*, *Necturus*) and those of the Caducibranchiata belonging to the family Derotremata (*Amphiuma*, *Menopoma*, *Cryptobranchus*).

It includes all (?) the papers on the embryology, nearly all those on the anatomy and histology. Most of the systematic, zoological and paleontological papers are also given.

#### A. Alphabetical arrangement of titles according to authors.

- 1873 ABBOTT, CHARLES C., Our largest Salamander (*Menopoma alleghaniensis*). *Science Gossip*, 1873, p. 271—272.
- 1878 ALBRECHT, PAUL, Ueber einen Processus odontoides des Atlas bei den urodelen Amphibien. *Centralbl. f. d. med. Wissensch.*, 1878, p. 578—579. Nachtrag, p. 705—706.
- 1903 ALLIS, E. P., On certain features of the cranial anatomy of *Bdellostoma dombeyi*. *Anat. Anz.*, Bd. 23, 1903, p. 259—281, 321—327.
- 1901 ANOEL, P., Étude du développement des glandes de la peau des Batraciens et en particulier de la Salamandre terrestre. *Arch. de Biol.*, T. 18, 1901, p. 257—289, 2 pl.
- 1892 ANDERSON, O. A., Zur Kenntnis des sympathischen Nervensystems der urodelen Amphibien. *Zool. Jahrb.*, Bd. 5, 1892, p. 184—210, 4 pl.
- 1896 ANDRES, ANGELO, La Salamandra gigantesca del Giappone (*Megalobatrachus maximus* BOUL.). *Atti. Soc. Ital. Sc. nat.*, Vol. 35, 1896, p. 201—218, 1 fig.
- 1898 ANDREWS, E. A., Filose activity in metazoan eggs. *Zool. Bull. Boston*, Vol. 2, 1898, p. 1—13, 5 fig.
- 1898a ANDREWS, E. A., Some activities of polar bodies. *Ann. Mag. Nat. Hist.*, Vol. 1, 1898, p. 109—116, 5 fig.
- 1849 BAIRD, SPENCER F., Revision of the North American tailed batrachians with descriptions of new genera and species. *Journ. Acad. Nat. Sci. Philad.*, Ser. 2, Vol. 1, 1849, p. 281—294.
- 1826 BARNES, DANIEL H., An arrangement of the genera of the Batrachian animals with a description of the more remarkable species including a monograph of the doubtful reptiles. *Amer. Journ. Sci. and Arts*, Ser. 1, Vol. 11, 1826, p. 268—297.
- 1827 BARNES, DANIEL H., Note on the doubtful reptiles. *Amer. Journ. Sci. and Arts*, Ser. 1, Vol. 13, 1827, p. 66—70.
- 1807 BARTON, BENJ. SMITH, (Scientific notes). *Philad. Med. and Physical Journ.*, Suppl. 2, 1807.
- 1808 BARTON, BENJ. SMITH, Notices of *Siren lacertina* and of another species of the same genus. Philadelphia 1808. 8°.
- 1812 BARTON, BENJ. SMITH, Some account of the *Siren lacertina* and other species of the same genus of amphibious animals. Philadelphia 1812. 8°.
- 1812a BARTON, BENJ. SMITH, A memoir concerning an animal of the class of Reptilia or Amphibia, which is known in the United States by the names of Alligator and Hellbender. Philadelphia 1812. 8°. 1 pl.
- 1885 BAUR, G., Einige Bemerkungen über die Ossifikation der „langen“ Knochen. *Zool. Anz.*, Bd. 8, 1885, p. 580—581.
- 1888 BAUR, G., Beiträge zur Morphologie des Carpus und Tarsus der Vertebraten. Teil 1. Batrachia. Jena 1888.
- 1889 BAUR, G., On the morphology of the vertebrate skull. 1. The otic elements. *Journ. Morphol.*, Vol. 3, 1889, p. 167—174.
- 1891 BAUR, G., The pelvis of the Testudinata, with notes on the evolution of the pelvis in general. *Journ. Morphol.*, Vol. 1, 1891, p. 345—359.
- 1894 BAWDEN, H. H., The nose and JACOBSON'S organ with especial reference to the Amphibia. *Journ. Comp. Neurol. and Psychol.*, Vol. 4, 1894, p. 117—152, 8 pl.
- 1878 BEALE, LIONEL S., The microscope in medicine. 4. ed. London 1878.
- 1799 BEAUVOIS, Memoir on a new species of *Siren* [operculata]. *Trans. Amer. Phil. Soc.*, Vol. 4, 1799, p. 277—281.
- 1902 BECKER, VICTOR, Untersuchungen an der Mundschleimhaut von *Cryptobranchus japonicus*. *Diss. Phil.*, Berlin 1902, p. 60.
- 1903 BEDDARD, F. E., Normally unequal growth as a possible cause of death. *Nature*, Vol. 68, 1903, p. 497.
- 1904 BEDDARD, F. E., On some points in the anatomy chiefly of the heart and vas-system of the Japanese Salamander, *Megalobatrachus japonicus*. *Proc. Zool. Soc. London*, Vol. 2, 1904, p. 298—315, 7 fig.
- 1899 BEER, TH., Die Accommodation des Auges bei den Amphibien. *Arch. Ges. Phys. PFLÜGER*, Bd. 73, 1899, p. 501—534, 14 Fig.

- 1835 BELL, THOMAS, Amphibia. In Todd's Cyclopaedia of anatomy and physiology, Vol. 1, London 1835/1836, p. 90—107.
- 1907 BENDER, OTTO, Die Homologie des Spritzloches der Selachier und der Paukenhöhle der Amphibien, Saurapsiden und Säugetiere auf Grund ihrer Innervation. Anat. Anz., Ergänzungsheft zu Bd. 30, 1907, p. 38—41.
- 1907a BENDER, OTTO, Die Schleimhautnerven des Facialis, Glossopharyngeus und Vagus. Studien zur Morphologie des Mittelohres und der benachbarten Kopfregion der Wirbeltiere. Denkschr. Med.-naturw. Ges. Jena, Bd. 7, 1907, p. 341—454.
- 1900 BENSLEY, R. R., The oesophageal glands of Urodela. Biol. Bull., Vol. 2, 1900, p. 87—101, 8 Fig.
- 1898 BERRY, JOHN M., A comparison of the phagocytic action of leucocytes in Amphibia and Mammalia. Trans. Amer. Micr. Soc., Vol. 19, 1898, p. 93—116, 5 pl.
- 1860 BETTZIECH, BETA H., Der Riesensalamander von Japan (*Megalobatrachus maximus*). Natur, Bd. 9, 1860, p. 211—212.
- 1846 BIDDER, F. H., Vergleichend-anatomische und histologische Untersuchungen über die männlichen Geschlechts- und Harnwerkzeuge der nackten Amphibien. Dorpat 1846. p. 1—71.
- 1889 BJELETZKIJ, N. F., Physiologische Bemerkung über den Riesensalamander (Russian). Arb. d. Naturf. Ges. Charkow, Bd. 15, 1889, p. 173—206.
- 1871 BLANCHARD, EMILE, On a new gigantic Salamander (*Sieboldia davidiana* BLANCH.) from Western China. Ann. Mag. Nat. Hist., Ser. 4, Vol. 8, 1871, p. 212—214.
- 1871a BLANCHARD, EMILE, Note sur une nouvelle Salamandre gigantesque (*Sieboldia davidiana* BLANCH.) de la Chine occidentale. Compt. Rend. Acad. Sc. Paris, 1871, p. 79—80.
- 1891 BLANCHARD, RAPH., Evacuation de noyaux cellulaires simulant une helminthiase et une coccidiose (*Proteus anguineus*). Bull. Soc. Zool. France, T. 16, 1891, p. 22—23.
- 1874 BLEYER, E., Magenepithel und Magendrüsen der Batrachier. Inaug.-Diss. Königsberg, 1874.
- 1881 BOAS, J. E. V., Ueber den conus arteriosus und die Arterienbogen der Amphibien. Morphol. Jahrb., Bd. 7, 1881, p. 488—572, 3 Taf.
- 1826 BOIE, H., Merkmale einiger japanischer Lurche. Isis, 1826, p. 203—216.
- 1899 BOLAU, HERM., Glandula thyroidea und Glandula Thymus der Amphibien. Zool. Jahrb., Abt. f. Anat., Bd. 12, 1899, p. 657—710, 11 Fig.
- 1832 BONAPARTE, CARLO LUCIANO, Fauna Italica. Roma 1832—1841, T. 2, Tab. 131.
- 1838 BONAPARTE, CARLO LUCIANO, Iconografia della Fauna Italica. Bibl. Ital., 93, 1838.
- 1839 BONAPARTE, CARLO LUCIANO, Amphibia europaea ad systema nostrorum vertebratorum ornitata. Turini 1839. 4°. p. 72.
- 1876 BÖTTGER, OSKAR, Ueber die äußeren Kiemenöffnungen bei jungen Exemplaren des japanischen Riesensmolchs. Zool. Garten, Bd. 17, 1876, p. 432—435.
- 1882 BOULENGER, G. A., Catalogue of the Batrachia gradientia s. caudata and Batrachia apoda in the Collection of the British Museum. 2. ed. London 1882. 8°. p. 8—127, 9 pl.
- 1907 BRACHET, A., Recherches sur l'ontogenèse de la tête chez les Amphibiens. Arch. de Biol., T. 23, 1907, p. 165—257.
- 1907 BRAUS, H., Ueber Frühanlagen der Schultermuskeln bei Amphibien und ihre allgemeinere Bedeutung. Anat. Anz., Erg.-Heft, Bd. 30, 1907, p. 192—219.
- 1900 BRIDGE, T. W., The air-bladder and its connection with the auditory organ in *Necturus borneensis*. Journ. Linn. Soc. Zool., Vol. 27, 1900, p. 503—540, 2 pl.
- 1903 BROILI, J. F., Zur Fortpflanzung des japanischen Riesensalamanders (*Megalobatrachus maximus*). Natur und Haus, Bd. 12, 1903.
- 1901 BROMAN, J., Notiz über das „Halsstück“ der Spermien von *Pelobates fuscus* nebst kritischen Bemerkungen über die Nomenklatur der Spermienchwanzfäden. Anat. Anz., Bd. 20, 1901, p. 347—351, 3 Fig.
- 1904 BROMAN, J., Die Entwicklungsgeschichte der Bursa omentalis und ähnliche Receptbildungen bei den Wirbeltieren. Wiesbaden 1904. p. 611, 650, Fig. 20 Pl.
- 1902 BUCHS, G., Ueber den Ursprung des Kopfskelettes bei *Necturus*. Leipzig 1902. 8°. Also Morph. Jahrb., Bd. 29, 1901, p. 582—613.
- 1873 BUGNION, E., Recherches sur les organes sensitifs qui se trouvent dans l'épiderme du Protée et de l'Axolotl. Bull. Soc. Vaudoise de Sci. nat. Lausanne, T. 12, 1873.
- 1874 BUGNION, E., Sur *Proteus anguineus*. Bull. Soc. Vaudoise Sci. nat. Lausanne, T. 12, 1874, p. 193—194.
- 1874a BUGNION, E., Sur le système nerveux du *Proteus anguineus*. Bull. Soc. Vaudoise Sci. nat. Lausanne, T. 13, 1874, p. 441—443.
- 1896 BUMPUS, H. C., A contribution to the study of variation. Skeletal variation of *Necturus maculatus* RAF. Journ. Morphol., Vol. 12, 1896, p. 455—484, 2 pl.
- 1904 BUSSY, L. P. DE, Eerste ontwikkelingsstadien van *Megalobatrachus maximus* SCHLEGEL. Amsterdam 1904.
- 1905 BUSSY, L. P. DE, Die ersten Entwicklungsstadien des *Megalobatrachus maximus* SCHLEGEL. Zool. Anz., Bd. 38, 1905.
- 1883 CAMERANO, LORENZO, Interno alla neotenia ed allo sviluppo degli Anfibi. Atti. Acc. Tor., T. 19, 1883, p. 84—93.
- 1883a CAMERANO, LORENZO, Ricerche intorno alla vita branchiale degli Anfibi. Zool. Anz., Bd. 6, 1883, p. 685—687.

- 1803 CAMPER, PETER, Du Dugon du comte de BUFFON et de la Sirena lacertina de chevalier LINNAEUS. Oeuvres., T. 3, Paris 1803.
- 1886 CARLSON, A. J., Untersuchungen über Gliedmaßenreste bei Schlangen. Bihang till K. Svenska Vet. Akad. Handlingar, Bd. 11, 1886.
- 1906 CARLSON, A. J., Die Ganglienzellen des Bulbus arteriosus und der Kammerspitze beim Salamander (*Necturus maculatus*). Arch. f. d. ges. Physiol., Bd. 109, 1906, p. 51—69, 3 Fig.
- 1879 CHAPMAN, HENRY C., Notes on the Amphiuma. Proc. Acad. Nat. Sci. Philadelphia 1879, p. 144—145.
- 1893 CHAPMAN, HENRY C., Observations on the Japanese Salamander *Cryptobranchus maximus* (SCHLEGEL). Proc. Acad. Nat. Sci. Philad., 1893, p. 227—233, 2 pl.
- 1883 CHAUVIN, MARIE VON, Vorläufige Mitteilung über die Fortpflanzung des *Proteus anguineus*. Zool. Anz., Bd. 6, 1883, p. 330—332.
- 1883a CHAUVIN, MARIE VON, Die Art der Fortpflanzung des *Proteus anguineus*. Zeitschr. wiss. Zool., Bd. 38, 1883, p. 671—685, 1 Taf.
- 1884 CHAUVIN, MARIE VON, Ueber die Färbung des Männchens von *Proteus anguineus*. Naturforscher, Bd. 16, 1884, p. 480.
- 1840 CHIAJE, STEFANO DELLE, Ricerche anatomico-biologiche sul Proteo serpentino. Napoli 1840. 21 p. 1<sup>o</sup>. 5 pl.
- 1897 CHORONSHITZKY, BORIS, Entstehung der Milz und des dorsalen Pankreas beim *Necturus*. Compt. Rend. XII. Congrès intern. méd. Moscow, Vol. 2, 1897, p. 115—120.
- 1900 CHORONSHITZKY, BORIS, Die Entstehung der Milz, Leber, Gallenblase, Bauchspeicheldrüse und des Pfortadersystems bei den verschiedenen Abteilungen der Wirbeltiere. Anat. Hefte, Bd. 13, 1900, p. 363—623, 85 Fig.
- 1882 CLAUS, C., Lehrbuch der Zoologie. 2 Bde. 4. ed. Leipzig 1882.
- 1893 CLAYPOLE, EDITH J., An investigation of the blood of *Necturus* and *Cryptobranchus*. Proc. Amer. Micr. Soc., Vol. 15, 1893, p. 39—76, 6 pl.
- 1896 CLAYPOLE, EDITH J., Notes on the comparative histology of blood and muscle. Trans. Amer. Micr. Soc., Vol. 18, 1896, p. 49—70, 5 pl.
- 1895 CLEMENS, P., Die äußeren Kiemen der Wirbeltiere. Anat. Hefte, Bd. 5, 1895, p. 51—156, 4 Taf., 5 Fig.
- 1821 CONFIGLIACHI, P., and RUSCONI, M., Observations on the natural history and structure of the *Proteus anguineus*. Edinb. Phil. Journ., Vol. 4, 1821, p. 388—406, and Vol. 5, p. 84—112.
- 1885 COPE, E. D., Amphibia. In Standard Natural History, Vol. 3, Boston 1885, p. 303—347.
- 1886 COPE, E. D., The retrograde metamorphosis of Siren. Amer. Nat., Vol. 19, 1886, p. 1226—1227.
- 1886a COPE, E. D., On the structure and affinities of the Amphiumidae. Amer. Philos. Soc., 1886.
- 1889 COPE, E. D., The Batrachia of North America. Bull. 34 U. S. Nat. Mus., Washington 1889.
- 1889a COPE, E. D., On the relations of the hyoid and otic elements of the skeleton in the Batrachia. Journ. Morphol., Vol. 2, 1889, p. 297—310, 3 pl.
- 1895 COPE, E. D., A careless writer on Amphiuma. Amer. Nat., Vol. 29, 1895, p. 1108—1110.
- 1860 CRISP, EDWARDS, Note on the blood-corpuscles of the Japanese gigantic Salamander (*Sieboldia maxima*). Proc. Zool. Soc. London, 1860, Part 28, p. 203—205.
- 1800 CUVIER, G., Sur le Siren lacertina. Bull. des. Sci. Soc. Philom., An. 8, T. 2, p. 106—107.
- 1827 CUVIER, G., Sur le genre de Reptiles batraciens nommé Amphiuma et sur une nouvelle espèce de cet genre (*A. tridactylum*). Men. du Muséum, T. 14, 1827, p. 1—14, 2 pl.
- 1831 CUVIER, G., Règne animale. Translated by EDWARD GRIFFITH and others, London 1831, Vol. 9, p. 412.
- 1853 DALTON, INO. C., Some account of the *Proteus anguineus*. Amer. Journ. Sci., Ser. 2, Vol. 15, 1853, p. 378—393.
- 1894 DAVISON, ALVIN, *Amphiuma tridactyla*. Princeton Coll. Bull., 1894.
- 1894a DAVISON, ALVIN, The arrangement of muscular fibers in *Amphiuma tridactyla*. Anat. Anz., Bd. 9, 1894, p. 332—336.
- 1895 DAVISON, ALVIN, A contribution to the anatomy and phylogeny of *Amphiuma means* (GARDNER). Journ. Morphol., Vol. 11, 1895, p. 375—410, 2 pl.
- 1896 DAVISON, ALVIN, The tentacular apparatus of *Amphiuma*. Amer. Nat., Vol. 30, 1896, p. 684—689, 4 fig.
- 1897 DAVISON, ALVIN, A preliminary contribution to the development of the vertebral column and its appendages. Anat. Anz., Bd. 14, 1897, p. 6—12, 7 Fig.
- 1834 DEEN, J. VAN, Over de zijdelingsche takken van de zwervende zenuw (*Nervus vagus*), van den *Proteus anguineus*. Tijdschr. voor natuurl. Geschied., Bd. 1, 1834, p. 112—129.
- 1842 DE KAY, JAMES E., Zoology of New York. Part 3. Reptiles and Amphibia. In Natural History of New York, Albany 1842. 4<sup>o</sup>.
- 1859 DOWLER, B., Natural history of the Amphiumidae with remarks on Crocodilian hibernation and instinct. New Orleans Med. and Surg. Journ., Vol. 16, 1859, p. 14—37.
- 1901 DRÜNER, L., Studien zur Anatomie der Zungenbein-, Kiemenbogen- und Kehlkopfmuskeln der Urodelen. 1. Thl. Zool. Jahrb., Bd. 15, 1901, p. 435—622, 7 Taf.
- 1903 DRÜNER, L., Ueber die Muskulatur des Visceralskelets der Urodelen. Anat. Anz., Bd. 23, 1903, p. 545—571, 16 Fig.

- 1903 DRZEWINA, ANNA, Sur le tissu lymphoïde du rein du *Proteus anguineus* LAUR. Note préliminaire. *Compt. Rend. Soc. Biol. Paris*, T. 55, 1903, p. 1091—1092.
- 1890 DUBOIS, RAFF., Sur la perception des radiations lumineuses par la peau chez les Protées aveugles des grottes de la Carniole. *Compt. Rend. Ac. Sc. Paris*, T. 110, 1890, p. 358—361.
- 1834 DUGES, A., Recherches sur l'ostéologie et la myologie des Batraciens. Paris 1831.
- 1831 DUMERIL, AND. MAR. CONST., et BIERON, G., *Erpétologie générale, ou histoire naturelle complète des Reptiles*. 8 Vol., Paris 1834—1844. 8°. Avec. pl.
- 1847 DUVERNOY, G. L., Ueber die Geschlechts- und die Harnorgane der dauerndgeschwänzten Batrachier. *Froniers Not.* 1847, No. 203.
- 1859 ECKER, A., Notiz über den Olm *Proteus anguinus* LAURENTI. *Corresp.-Bl. d. Naturf. Ver. Riga*, Bd. 11, 1859, p. 123—124.
- 1902 ECKEL, E. C., and PAULMIRE, F. C., Catalogue of the Reptiles and Batrachians New York State. *N. Y. State Museum Bull.*, Vol. 10, 1902, No. 51, p. 353—358, 410—414, 1 pl.
- 1904 EDINGER, LUDWIG, Vorlesungen über den Bau der nervösen Zentralorgane des Menschen und der Tiere. 2 Bde. 7. ed. Leipzig 1904, 1908.
- 1862 EHRENBERG, CHR., Ueber die Nahrung des *Proteus anguinus*. *Sitzungsber. d. Ges. naturf. Freunde Berlin*, 1862.
- 1867 EHRENBERG, CHR., Ueber den seit 7 Jahren lebend gehaltenen *Proteus anguinus*. *Sitzungsber. d. Ges. naturf. Freunde Berlin*, 1867, p. 1.
- 1868 EHRENBERG, CHR., Ueber den fast 9 Jahre in der Gefangenschaft gehaltenen *Proteus anguinus*. *Sitzungsber. d. Ges. naturf. Freunde Berlin*, 1868, p. 14—15.
- 1870 EHRENBERG, CHR., Ueber den in Gefangenschaft gehaltenen *Proteus anguinus*. *Sitzungsber. d. Ges. naturf. Freunde Berlin*, 1870, p. 2.
- 1872 EHRENBERG, CHR., Ueber den seit 1859 lebend gehaltenen *Proteus anguinus*, und über den seit 1860 beobachteten *Triton lacustris*. *Sitzungsber. d. Ges. naturf. Freunde Berlin*, 1872, p. 17—18.
- 1874 EHRENBERG, CHR., Ueber den seit 1859 lebend erhaltenen *Proteus anguinus*. *Sitzungsber. d. Ges. naturf. Freunde Berlin*, 1874, p. 9—10.
- 1899 EIGENMANN, C. H., A case of convergence. *Science*, Vol. 9, 1899, p. 280—282, 3 fig.
- 1900 EIGENMANN, C. H., The eyes of the blind vertebrates of North America. 2. The eyes of *Typhlomolge rathbuni* STEJ. *Trans. Amer. Micr. Soc.*, Vol. 21, 1900, p. 49—60.
- 1900a EIGENMANN, C. H., Degeneration in the eyes of cold-blooded vertebrates of the North American caves. *Science*, Vol. 11, 1900, p. 492—503, 14 fig.
- 1900 EIGENMANN, C. H., and DENNY, W. A., Eyes of blind vertebrates of North America. 3. The structure and ontogenic degeneration of the eyes of the Missouri Cave Salamander. *Biol. Bull.*, Vol. 2, 1900, p. 33—41, 9 fig.
- 1900a EIGENMANN, C. H., and DENNY, W. A., Eyes of Cave Salamander (*Typhlotriton spelaeus*). *Proc. Indiana Acad. Sci.*, 1898, p. 252—255.
- 1899 EISEN, GUSTAV, Blood plates of the human blood with notes on the erythrocytes of *Amphiuma* and *Necturus*. *Journ. Morphol.*, Vol. 15, 1899, p. 635.
- 1875 EISMANN, GUST., Der Adelsberger Olm *Proteus anguinus* in Gefangenschaft. *Zool. Garten*, Bd. 16, 1875, p. 394—396.
- 1766 ELLIS, JOHN, An account of an Amphibious Bipes (*Siren lacertina*). *Phil. Trans. Roy. Soc. London*, Vol. 56, 1766, p. 189—192, 1 pl.
- 1905 EMERSON, E. T., General anatomy of *Typhlomolge rathbuni*. *Proc. Boston Soc. Nat. Hist.*, Vol. 32, 1905, p. 43—76, pl. 1.
- 1897 EMERY, C., Die fossilen Reste von *Archegosaurus* und *Eryops* und ihre Bedeutung für die Morphologie des Gliedmaßenskelets. *Anat. Anz.*, Bd. 14, 1897, p. 201—208, 7 Fig.
- 1863 ERBER, JOSEPH, Beobachtungen an Amphibien in der Gefangenschaft. *Verh. d. Zool.-bot. Ges. Wien*, Bd. 13, 1863, p. 129—132.
- 1864 ERBER, JOSEPH, Die Amphibien der österr. Monarchie. Mit Anführung der Beobachtungen die an den in der Gefangenschaft gehaltenen Arten gemacht wurden. *Verh. d. zool. bot. Ges. Wien*, Bd. 14, 1864, p. 697—720.
- 1877 ERBER, JOSEPH, Ueber die Lebensweise von *Siren lacertina* in der Gefangenschaft. *Verh. d. Zool.-bot. Ges. Wien*, Bd. 26, 1877, p. 114—116.
- 1893 EYCLESYMER, ALBERT C., Development of the optic vesicles in Amphibia. *Journ. Morphol.*, Vol. 8, 1893, p. 189—194.
- 1895 EYCLESYMER, ALBERT C., Early development of *Amblystoma*. *Journ. Morphol.*, Vol. 10, 1895, p. 345—400.
- 1902 EYCLESYMER, ALBERT C., The formation of the embryo of *Necturus*, with remarks on the theory of conerescence. *Anat. Anz.*, Bd. 21, 1902, p. 341—353.
- 1902a EYCLESYMER, ALBERT C., Nuclear changes in the striated muscle cell of *Necturus*. *Anat. Anz.*, Bd. 21, 1902, p. 379—385.
- 1904 EYCLESYMER, ALBERT C., The cytoplasmic and nuclear changes in the striated muscle cell of *Necturus*. *Amer. Journ. Anat.*, Vol. 3, 1904, p. 285—310, 4 pl.

- 1904a EYCLESHYMER, ALBERT C., Bilateral symmetry in the egg of *Necturus*. *Anat. Anz.*, Bd. 25, 1904, p. 230—240, 47 Fig.
- 1906 EYCLESHYMER, ALBERT C., The habits of *Necturus maculosus*. *Amer. Nat.*, Vol. 40, 1906, p. 123—136.
- 1906a EYCLESHYMER, ALBERT C., The growth and regeneration of the gills in the young *Necturus*. *Biol. Bull.*, Vol. 10, 1906, p. 171—175, 1 fig.
- 1906b EYCLESHYMER, ALBERT C., The development of the chromatophores in *Necturus*. *Amer. Journ. Anat.*, Vol. 5, 1906, p. 309—330, 7 fig.
- 1907 EYCLESHYMER, ALBERT, The Closing of Wounds in the Larval *Necturus*. *Amer. Journ. of Anat.*, Vol. VII, p. 317—326.
- 1906 FAVARO, GIUS., Ricerche anatomo-embriologiche intorno alla circolazione caudale ed ai cuori linfatici posteriori degli Anfibi, con particolare riguardo agli Urodeli. *Atti Accad. Sc. Venet. Trent. Padova*, Anno 3, 1906, p. 122—166, 20 fig.
- 1906 FELIX, W., and BUHLER, A., Die Entwicklung der Harn- und Geschlechtsorgane. *Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG*, Jena 1906, Bd. 3, p. 81.
- 1894 FIELD, HERBERT H., Sur le développement des organes excréteurs chez l'Amphiuma. *Compt. Rend. de l'Acad. Sci. Paris*, T. 118, 1894, p. 1221—1224.
- 1895 FIELD, HERBERT H., Bemerkungen über die Entwicklung der Wirbelsäule bei den Amphibien; nebst Schilderung eines abnormen Wirbelsegmentes. *Morphol. Jahrb.*, Bd. 22, 1895, p. 340—356.
- 1843 FISCHER, J. G., *Amphibiorum nuderum neurologia*. Berolini 1843.
- 1864 FISCHER, J. G., *Anatomische Abhandlungen über die Perennibranchiaten und Derotremen*. Hamburg 1864. 4<sup>o</sup>. p. 170, 6 pl.
- 1900 FISCHER, SIGWART, Ueber *Proteus anguineus* L. *Verh. Schweiz. Naturf. Ges.*, 1900, p. 76.
- 1826 FITZINGER, LEOPOLD J., *Neue Klassifikation der Reptilien*. Wien 1826. 4<sup>o</sup>.
- 1850 FITZINGER, LEOPOLD J., Ueber den *Proteus anguineus*. *Wien. Sitzungsber., Math.-nat. Kl.*, Bd. 5, 1850, p. 291—303.
- 1909 FLEISSIG, JULIUS, Zur Anatomie der Nasenhöhle von *Cryptobranchus japonicus*. *Anat. Anz.*, Bd. 35, 1909, p. 48—54, 5 Fig.
- 1861 FRAUENFELD, G. Ritter von, Notiz über den Olm. *Verh. d. Zool.-bot. Ges. Wien*, Bd. 12, 1861, p. 29.
- 1882 FREAR, WM., Vitality of the Mud Puppy (*Menopoma*). *Amer. Nat.*, Vol. 16, 1882, p. 325—326.
- 1846 FREYER, HEINRICH, Ueber eine neue Art von *Hypochthon* (*Proteus*). *Arch. f. Naturgesch.*, Bd. 12, 1846, p. 289—290.
- 1894 FÜLLEBORN, F., Bericht über eine zur Untersuchung der Entwicklung von *Amia*, *Lepidosteus* und *Necturus* unternommene Reise nach Nordamerika. *Sitzungsber. d. Akad. d. Wiss. Berlin*, Bd. 40, 1894, p. 1057—1070.
- 1877 FÜRBRINGER, M., Zur Entwicklung der Amphibienniere. Heidelberg 1877.
- 1906 FÜRBRINGER, M., Zur vergleichenden Anatomie des Brustschulterapparates und der Schultermuskeln. *Jena. Zeitschr.*, Bd. 36, 1902, p. 289—736, 5 Taf., 23 Fig.
- 1895 GADOW, HANS, On the evolution of the vertebral column of Amphibia and Amniota. *Proc. Roy. Soc. London*, Vol. 58, 1895, p. 257—259.
- 1901 GADOW, HANS, *Amphibia and Reptilia*. Cambridge Natural History, Vol. 8, London 1901.
- 1882 GAGE, SIMON HENRY, Observations on the fat cells and connective tissue corpuscles of *Necturus* (*Menobranchus*). *Proc. Amer. Soc. Micr. Buffalo*, Vol. 5, 1882, p. 109—126, 1 pl.
- 1885 GAGE, SIMON HENRY, Notes on the epithelium lining the mouth of *Necturus* and *Menopomia*. *Proc. Amer. Soc. Micr. Buffalo*, Vol. 8, 1885, p. 126.
- 1885a GAGE, SIMON HENRY, Notes on the blood corpuscles of *Necturus*. *Proc. Amer. Soc. Micr. Buffalo*, Vol. 8, 1885, p. 126.
- 1890 GAGE, SUSANNA PHELPS, The intramuscular endings of fibres in the skeletal muscles of the domestic and laboratory animals. *Proc. Amer. Soc. Micr.*, Vol. 13, 1890, p. 132—138, 1 pl.
- 1890 GAGE, SIMON HENRY, and GAGE, SUSANNA PHELPS, Changes in the ciliated areas of the alimentary canal of the alimentary canal of the amphibia and the relation to the mode of respiration. *Proc. Amer. Assoc. Adv. Sci.*, Vol. 39, 1890, p. 337—338.
- 1897 GALEOTTI, GINO, Studio morfologico e citologico della volta del diencefalo in alcuni Vertebrati. *Riv. Pat. nerv. e ment.* Firenze, Vol. 2, 1897, p. 481—517, 20 fig.
- 1888 GARNIER, J. H., On a new species of *Menobranchus* (*lateralis* var. *Latastei*). *Proc. Canad. Inst., Ser. 3*, Vol. 5, 1888, p. 218—219.
- 1901 GAUFF, ERNST, Das Hypobranchialskelet der Wirbeltiere. *Ergeb. Anat. u. Entwickl.*, Bd. 14, 1904, p. 808—1048, 46 Fig.
- 1906 GAUFF, ERNST, Die Entwicklung des Kopfskeletes. *Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG*. Bd. 3, Jena 1906.
- 1883 GEERTS, A. J. C., Notice sur la grande Salamandre du Japon. *Cryptobranchus japonicus* v. HOEVEN. *Nouv. Arch. Mus. Paris*, T. 2, 1883, p. 273—290, 1 pl.
- 1881 GEERTS, A. J. C., Notice sur la grande Salamandre du Japon. *Nouv. Arch. du Mus.*, T. 5, 1884, p. 273—290.
- 1862 GEGENBAUER, CARL, Untersuchungen zur vergleichenden Anatomie der Wirbelsäule bei Amphibien und Reptilien. Leipzig 1862. 4<sup>o</sup>. p. 22.

- 1898 GEGENBAUR, CARL, Vergleichende Anatomie der Wirbeltiere. 2 Bde. Leipzig 1898.
- 1905 GEYER, HANS, Der Schlammteufel (*Cryptobranchus alleghaniensis*). Natur u. Haus, Bd. 13, 1905, p. 356—358, 1 Fig.
- 1850 GIBBES, LEWIS R., On a new species of *Menobranchus* from South Carolina (*M. punctatus*). Proc. Amer. Assoc. Adv. Sci., Vol. 3, 1850, p. 159.
- 1853 GIBBES, LEWIS R., Description of *Menobranchus punctatus*. Boston Journ. Nat. Hist., Vol. 6, 1853, p. 369—573, Fig. 1.
- 1806 GMELIN, CARL CHRISTL, Gemeinnützliche systematische Naturgeschichte für gebildete Leser nach dem LINNÉschen Natursystem. (German, Latin and French text.) 4 Teile. Mannheim 1806—1829.
- 1891 GÖPPERT, E., Die Entwicklung und das spätere Verhalten des Pancreas der Amphibien. Morphol. Jahrb., Bd. 17, 1891, p. 100—122, 1 Taf.
- 1894 GÖPPERT, E., Die Kehlkopfmuskulatur der Amphibien. Morphol. Jahrb., Bd. 22, 1894, p. 1—78.
- 1895 GÖPPERT, E., Zur Kenntnis der Amphibienrippen. Morphol. Jahrb., Bd. 22, 1895, p. 411—448, 5 Fig.
- 1896 GÖPPERT, E., Die Morphologie der Amphibienrippen. Festschr. f. GEGENBAUR, Bd. 1, 1896, p. 393—435, 2 Taf., 10 Fig.
- 1898 GÖPPERT, E., Der Kehlkopf der Amphibien und Reptilien. Morphol. Jahrb., Bd. 26, 1898, p. 282, 2 Taf., 5 Fig.
- 1906 GÖPPERT, E., Die Entwicklung des Mundes und der Mundhöhle mit Drüsen und Zunge; die Entwicklung der Schwimmblase, der Lunge und des Kehlkopfes der Wirbeltiere. Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG, Jena 1906, Bd. 2, p. 1—108.
- 1825 GRAY, JOHN EDWARD, A synopsis of the genera of Reptiles and Amphibia, with a description of some new species. Thomson Ann. Philos., Vol. 10, 1825, p. 193—217.
- 1857 GRAY, JOHN EDWARD, On the genus *Necturus* or *Menobranchus* with an account of its skull and teeth. Proc. Zool. Soc. London, Vol. 25, 1857, p. 61—64.
- 1873 GRAY, JOHN EDWARD, On a Salamander (*Sieboldia*) from Shanghai. Ann. Mag. Nat. Hist., Ser. 4, Vol. 12, 1873, p. 188.
- 1896 GREEN, ISABELLA M., The peritoneal epithelium in Amphibia. Amer. Nat., Vol. 30, 1896, p. 944—945.
- 1897 GREEN, ISABELLA M., The peritoneal epithelium of some Ithaea Amphibia (*Necturus*, *Amyblystoma*, *Desmognathus* and *Diemyctylus*). Trans. Amer. Micr. Soc., Vol. 18, 1897, p. 76—106, 5 pl.
- 1876 GROTE, A. R., On casting the skin in *Menopoma* (*Cryptobranchus*) *alleghaniensis*. Amer. Journ. Sci., Ser. 3, Vol. 12, 1876, p. 472.
- 1876a GROTE, A. R., Note on *Menopoma alleghaniensis* of Harlan. Amer. Journ. Sci., Vol. 12, 1876.
- 1866 GRUBE, E., Ueber die Nahrung des *Proteus anguinus*. Jahrb. d. Schles. Ges. f. vaterl. Kultur, 1866, p. 63—64.
- 1896 GUERNE, JULES DE, Mort de la grande Salamandre du Japon (du Muséum d'Histoire naturelle de Paris). Etangs et Rivières, Vol. 10, 1896, p. 207.
- 1873 GULLIVER, GEORGE, Measurements of the red blood corpuscles of Batrachians. Proc. Zool. Soc. London, 1873, p. 162—165.
- 1904 HALL, R. W., The development of the mesonephros and the Müllerian ducts in Amphibia. Bull. Mus. Comp. Zool. Harvard, Vol. 45, 1904, p. 31—125, 8 pl., 1 fig.
- 1856 HALLOWELL, EDW., *Pseudotriton marginatus* and *flavissimus* nn. spp. Urodel. from Georgia. Proc. Acad. Nat. Sci. Philad., Vol. 8, 1856, p. 130—131.
- 1858 HALLOWELL, EDW., On the caducibranchiate Urodele Batrachians. Journ. Acad. Nat. Sci. Philad., Ser. 2, Vol. 3, 1858, p. 337.
- 1892 HARGITT, CHARLES W., On some habits of *Amphiuma means*. Science, Vol. 20, 1892, p. 159.
- 1823 HARLAN, RICHARD, Description of a Batrachian animal in a living state (*Amphiuma means*). Journ. Acad. Nat. Sci. Philad., Vol. 3, 1823, p. 54—59.
- 1824 HARLAN, RICHARD, Observations on the genus *Salamandra* with the anatomy of the *Salamandra gigantea* (BARTON) or *S. alleghaniensis* (MICHAUX) and two new genera proposed. Ann. Lye. Nat. Hist., Vol. 1, 1824, Pt. 2, p. 222—234.
- 1824a HARLAN, RICHARD, Dissection of a Batrachian animal in a living state (*Amphiuma means*). Philos. Mag., Vol. 63, 1824, p. 325—328, 1 fig.
- 1826 HARLAN, RICHARD, Ueber *Amphiuma means*. FORST. Not., Bd. 14, 1826, p. 8—10, Fig.
- 1826a HARLAN, RICHARD, Further observations on the *Amphiuma means*. Ann. Lye. Nat. Hist. New York, Vol. 1, 1826, p. 269.
- 1827 HARLAN, RICHARD, American herpetology, or Genera of the North American Reptilia; with a synopsis of the species. Journ. Acad. Nat. Sci. Philad., Vol. 5, 1827, Pt. 2, p. 324.
- 1835 HARLAN, RICHARD, Medical and physical researches; or Original memoirs. Philad. 1835, p. 164—176, fig.
- 1858 HARTING, PIETER, Note sur les corpuscles sanguins du *Cryptobranchus japonicus*. Verslagen en Mededeelingen der Koninklijke Akad. van Wetensch., Afdeling Natuurkunde, Vol. 7, 1858, p. 368—372, 1 pl.
- 1871 HARTING, PIETER, Nieuwe Reuzen-Salamander. Album der Natur, 1871.
- 1888 HAY, O. P., Observations on *Amphiuma* and its young. Amer. Nat., Vol. 22, 1888, p. 315—321.
- 1889 HAY, O. P., On the structure of the skull of the larva of *Amphiuma*. Proc. Amer. Assoc. Adv. Sci., 1889, p. 286.
- 1890 HAY, O. P., The skeletal anatomy of *Amphiuma* during its earlier stages. Journ. Morphol., Vol. 4, 1890, p. 11—34, 1 pl.

- 1892 HAY, O. P., The Batrachians and Reptiles of Indiana. Indianapolis 1892. 8°. p. 204, 3 pl.
- 1893 HEIDENHAIN, M., Die Hautdrüsen der Amphibien. Sitz.-Ber. Physik.-med. Ges. Würzburg, 1893, No. 4, p. 52—64.
- 1900 HEIDENHAIN, M., Ueber die Zentralkapseln und Pseudochromosomen in den Samenzellen von Proteus, sowie über ihr Verhältnis zu den Idiosomen, Chondromiten und Archoplasmenschleifen. Anat. Anz., Bd. 18, 1900, p. 513—550, 8 Fig.
- 1907 HEIDENHAIN, M., Plasma und Zelle. Handbuch der Anatomie des Menschen von KARL VON BARDELEBEN, Bd. 8, Abt. 2, Jena 1907.
- 1907 HERRNUN, CLARA, A peculiar pelvic attachment in *Necturus maculatus*. Biol. Bull., Vol. 12, 1907, p. 375—377.
- 1893 HERRICK, C. L., The callosum and hippocampal region in marsupial and lower brains. Journ. Comp. Neurol. and Psychol., Vol. 3, 1893, p. 176—182.
- 1894 HERRICK, C. JUDSON, The cranial nerves of *Amblystoma*. Journ. Comp. Neurol. and Psychol., Vol. 4, 1894, p. 193—207.
- 1909 HERRICK, C. JUDSON, The nervus terminalis (nerve of PINKUS) in the Frog. Journ. Comp. Neurol. and Psychol., Vol. 19, 1909, p. 203—209, 1 Fig.
- 1906 HERTWIG, OSCAR, Allgemeine Biologie. Jena 1906. 8°.
- 1889 HESS, C., Beschreibung des Auges von *Talpa europaea* und von *Proteus anguineus*. Arch. f. Ophthalmol., Bd. 35, 1889, p. 1—19, 1 Taf.
- 1909 HIRSCH-TABOR, O., Ueber das Gehirn von *Proteus anguineus*. Arch. mikr. Anat. u. Entw., Bd. 72, 1909, p. 719—731.
- 1906 HOCHSTETTER, F., Die Entwicklung des Blutgefäßsystems. Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG, Jena 1906, Bd. 3, p. 21—166.
- 1838 HOEVEN, J. VAN DER, [Letter to Mr. OGILBY]. Proc. Zool. Soc. London, 1838, p. 25.
- 1838a HOEVEN, J. VAN DER, Jets over den grooten zoogenoemden Salamander van Japan. Met afbeeldingen van schedels en eene nieuwe afteekening van de *Menopoma* van Harlan. Tijdschr. Natuurl. Geschied. Physiol., Vol. 4, 1838, p. 375—386, 2 pl.
- 1838b HOEVEN, J. VAN DER, Sur une nouvelle espèce de *Cryptobranchus* du Japon. Bull. Sci. phys. natur. Néerlande, Leiden 1838, p. 90—91.
- 1839 HOEVEN, J. VAN DER, Note sur une nouvelle espèce de *Cryptobranchus*. Ann. Sci. nat., T. 11, 1839, p. 63—64.
- 1841 HOEVEN, J. VAN DER, Groote bloedschijljes bij *Cryptobranchus japonicus*. Tijdschr. voor Natuurl. Geschied., Vol. 8, 1841, p. 270—272.
- 1866 HOEVEN, J. VAN DER, Note sur le carpe et le tarse du *Cryptobranchus japonicus*. Arch. néerl. Sc. exact. et nat., T. 1, 1866, p. 321—327.
- 1866a HOEVEN, J. VAN DER, Considérations sur le genre *Menobranchus* et sur ses affinités naturelles. Arch. néerl. Sc. exact. et nat., T. 1, 1866, p. 1—16.
- 1866a HOEVEN, J. VAN DER, Notes on the genus *Menobranchus* and its natural affinities. Ann. Mag. Nat. Hist., Vol. 18, 1866, p. 363—375.
- 1867 HOEVEN, J. VAN DER, Ontleed en Dierkundige Bijdragen tot de Kennis van *Menobranchus*. Arch. néerl. Sc. exact. et nat., T. 2, 1867, p. 288.
- 1867a HOEVEN, J. VAN DER, Les globules du sang du *Menobranchus*. Arch. néerl. Sc. exact. et nat., T. 2, 1867, p. 288.
- 1867b HOEVEN, J. VAN DER, Ontleed en dierkundige bijdragen tot de Kennis der *Menobranchus*, den *Proteus* der meren van Noord-Amerika. Leiden 1867. 3 pl.
- 1867c HOEVEN, J. VAN DER, Over *Menobranchus*. Proc. verb. Vergad. Kon. Akad. Wetensch. Amsterdam, Afd. Natuurk., 1867, p. 3.
- 1877 HOFFMANN, C. K., Amphibien. BRONN'S Klassen und Ordnungen des Tierreiches, Leipzig 1877, Bd. 6, Abt. 2.
- 1902 HOFFMANN, C. K., Zur Entwicklungsgeschichte des Sympathicus. 2. Die Entwicklungsgeschichte des Sympathicus bei den Urodelen. Verh. Akad. Amsterdam, Vol. 8, 1902, No. 3, p. 101, 4 pl.
- 1836 HOLBROOK, JOHN EDWARDS, North American Herpetology: or a Description of the Reptiles inhabiting the United States. Vol. 5, Col. Pl. Philad., 1836—43. roy. 4°.
- 1901 HOLMGREN, E., Zur Kenntnis der cylindrischen Epithelzellen. Arch. mikr. Anat. u. Entw., Bd. 65, 1903, p. 280—297, 2 Taf., 5 Fig.
- 1903 HOUGHTON, HENRY S., Muscular and skeletal elements in *Spelerpes longicaudus*. Ohio Naturalist, Vol. 3, 1903, p. 379—393, fig. 1.
- 1908 HOWARD, ARTHUR DAY, The visual cells in vertebrates, chiefly in *Necturus*. Journ. Morphol., Vol. 19, 1908, p. 561—629, 5 pl.
- 1888 HOWES, G. B., and DAVIES, A. M., Observations upon the morphology and genesis of supernumerary phalanges with especial references to those of Amphibia. Proc. Zool. Soc. London, 1888.
- 1893 HOWES, G. B., Notes on variations and development of the vertebral and limb skeleton of the Amphibia. Proc. Zool. Soc. London, 1893, Pt. 2, p. 268—278, 15 fig.
- 1805 HUMBOLDT, ALEXANDER VON, Recueil d'observations de zoologie et d'anatomie comparée, 2 T., Paris 1805—32, 55 pl.

- 1872 HUMPHRY, G. M., The muscles and nerves of the *Cryptobranchus japonicus*. Journ. Anat. and Physiol., Vol. 6, 1872, p. 1—61, 4 pl.
- 1766 HUNTER, JOHN, Supplement to [JOHN ELLIS'S] account etc. being the anatomic description of the said animal [*Siren lacertina*]. Phil. Trans. Roy. Soc. London, Vol. 56, 1766, p. 307—310.
- 1874 HUXLEY, THOMAS HENRY, On the structure of the skull and heart of *Menobranchus*. Proc. Zool. Soc. London, 1874, p. 186—204, 4 pl.
- 1875 HUXLEY, THOMAS HENRY, Amphibia. In: Encyclopaedia Britannica, 9. ed., Vol. 1, 1875, p. 750—771.
- 1850 HYRTL, J., Bemerkungen über den *Proteus anguinus*. Wien, Sitz.-Ber., math.-nat. Kl., Bd. 5, 1850, p. 303.
- 1865 HYRTL, J., *Cryptobranchus japonicus*. Schediasma anatomicum. Vindobonae 1865.
- 1865a HYRTL, J., Kurze Inhaltsanzeige einer im nächsten Jahr zu veröffentlichenden Abhandlung über die Anatomie des Riesensalamanders. Wien, Sitz.-Ber., math.-nat. Kl., Bd. 1, 1865, p. 48—49.
- 1900 ISHIKAWA, C., Zur Entwicklungsgeschichte von *Cryptobranchus japonicus*. Toga Gakuge-Zasshi, 1900.
- 1902 ISHIKAWA, C., Ueber den Riesensalamander Japans. Mitt. Ges. f. Naturk. Ostasiens, Bd. 9, 1902, p. 79—94.
- 1903 ISHIKAWA, C., Distribution, oviposition, habits etc. of the Japanese giant-salamander [Japanese]. Tokyo Teishitsu Hakubutsukwan, Vol. 2, 1903, p. 27.
- 1904 ISHIKAWA, C., Beiträge zur Kenntnis des Riesensalamanders (*Megalobatrachus maximus* SCHL.). Proc. Depart. Nat. Hist. Tokyo, Imperial Museum, Vol. 1, 1904.
- 1905 ISHIKAWA, C., The gastrulation of the gigantic Salamander *Megalobatrachus maximus*. Zool. Magazine, Vol. 17, 1905.
- 1908 ISHIKAWA, C., Ueber den Riesensalamander Japans (*Cryptobranchus japonicus*). Entwicklung der äußeren Körperform. Mitt. Ges. f. Naturk. Ostasiens, 1908, 22 p., 3 Taf., 3 Fig.
- 1823 JAMES, EDWIN, Account of an expedition from Pittsburg to the Rocky Mountainis 1819. From the notes of Major LONG, Mr. T. SAY and others. 2 Vols. Philadelphia 1823. 8°. With atlas.
- 1897 JELLIFFE, SMITH ELY, Preliminary note upon the cytology of the brains of some Amphibians. I. *Necturus*. Journ. Comp. Neurol. and Psychol., Vol. 7, 1897, p. 146—151.
- 1876 JOHNSTON, CHRISTOPHER, Observations upon spermatozoa of *Amphiuma tridactylum*. Monthly Micr. Journ., Vol. 16, 1876, p. 61—63, 1 pl.
- 1903 JOHNSTON, J. B., Movements of the cerebro-spinal fluid in *Cryptobranchus*. Science, N. S., Vol. 17, 1903, p. 530.
- 1906 JOHNSTON, J. B., The nervous system of Vertebrates. Philadelphia 1906.
- 1905 JOSEPH, H., [Randreifen der roten Blutkörperchen von *Proteus*]. Verh. Anat. Ges. 19. Vers., 1905, p. 229—230.
- 1905a JOSEPH, H., Ueber die Zentralkörper der Nierenzelle. Verh. Anat. Ges. 19. Vers., 1905, p. 178—187, 16 Fig.
- 1905 KAMMERER, P., Ueber die Abhängigkeit des Regenerationsvermögens der Amphibienlarven von Alter, Entwicklungsstadium und spezifischer Größe. Arch. Entw.-Mech., Bd. 19, 1905, p. 148—180.
- 1905a KAMMERER, P., Die angebliche Ausnahme von der Regenerationsfähigkeit bei den Amphibien. Centralbl. Physiol., Bd. 19, 1905, p. 684—687.
- 1907 KAMMERER, P., Die Fortpflanzung des Grottenolmes (*Proteus anguineus* LAURENTI). Verh. k. k. Zool.-bot. Ges. Wien, Bd. 57, 1907, p. 277—292.
- 1903 KAMPEN, P. N. VAN, Eieren en spermien van *Megalobatrachus maximus* SCHL. Tijdsch. Ned. Dierk. Ver., Vol. 8, 1903.
- 1901 KEIBEL, F., Die Gastrulation und die Keimblattbildung der Wirbeltiere. Ergeb. Anat. Entw., Bd. 10, 1901, p. 1002—1119, 43 Fig.
- 1906 KEIBEL, F., Die Entwicklung der äußeren Körperform der Wirbeltierembryonen, insbesondere der menschlichen Embryonen aus den ersten 2 Monaten. Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG. Bd. 1, 1906, p. 174.
- 1903 KERBERT, C., Eieren van *Megalobatrachus maximus* SCHL. Tijdschr. d. Ned. Dierk. Ver., Vol. 8, 1903.
- 1904 KERBERT, C., Zur Fortpflanzung von *Megalobatrachus maximus* SCHL. Zool. Anz., Bd. 27, 1904, p. 305—320, 1 Fig.
- 1905 KERBERT, C., Ueber die Eier und Larven von *Megalobatrachus maximus* SCHL. Compt. Rend. 6 Cong. Internat. Zool. Bâle, 1905, p. 289—294.
- 1894 KINGSBURY, B. F., The histological structure of the enteron of *Necturus maculatus*. Proc. Amer. Micr. Soc. Ithaca, Vol. 16, 1894, p. 19—64, 8 pl.
- 1895 KINGSBURY, B. F., The spermatheca and methods of fertilization in some American Newts and Salamanders. Trans. Amer. Micr. Soc., Vol. 17, 1895, p. 261—304, 4 pl.
- 1895a KINGSBURY, B. F., On the brain of *Necturus maculatus*. Journ. Comp. Neurol. and Psychol., Vol. 5, 1895, p. 139—205, 3 pl.
- 1895b KINGSBURY, B. F., The lateral line system of sense organs in some American Amphibia and comparisons with Dipnoans. Proc. Amer. Micr. Soc., Vol. 17, 1895, p. 115—150, 5 pl.
- 1903 KINGSBURY, B. F., Columella auris and Nervus facialis in the Urodela. Journ. Comp. Neurol. and Psychol., Vol. 3, 1903, p. 313—334.
- 1904 KINGSBURY, B. F., The rank of *Necturus* among tailed Batrachia. Biol. Bull., Vol. 8, 1904, p. 67—74.

- 1908 KINGSBURY, B. F., and REED, H. D., The columella aurita in Amphibia. *Anat. Record.*, Vol. 2, 1908, p. 81—91.
- 1892 KINGSLEY, J. S., The head of an embryo Amphiuma. *Amer. Nat.*, Vol. 26, 1892, p. 671—680.
- 1899 KINGSLEY, J. S., and RUDDICK, W. H., The ossicula auditus and mammalian ancestry. *Amer. Nat.*, Vol. 33, 1899, p. 219—230, 3 fig.
- 1900 KINGSLEY, J. S., The ossicula auditus. *Tufts Coll. Stud.*, 1900, No. 6, p. 203—274, with pl.
- 1902 KINGSLEY, J. S., The systematic position of the Caecilians. *Tufts Coll. Stud.*, 1902, No. 7, p. 293—321, 3 pl.
- 1883 KLAUSSNER, F., Das Rückenmark des *Proteus anguineus* L. München 1883.
- 1886 KNAPE, E., Das BIDDERSche Organ. *Morphol. Jahrb.*, Bd. 11, 1886, p. 489—552.
- 1856 KNEELAND, SAMUEL, Letter relating to a supposed new species of *Siredon* from Lake Superior. *Proc. Boston Soc. Nat. Hist.*, Vol. 6, 1856—59, p. 152—154.
- 1856a KNEELAND, SAMUEL, On the breathing apparatus of the *Menobranchus*. *Proc. Boston Soc. Nat. Hist.*, Vol. 6, 1856—59, p. 428—430.
- 1858 KNEELAND, SAMUEL, [Notes on *Menobranchus*]. *Proc. Boston Soc. Nat. Hist.*, Vol. 6, 1858, p. 152—154, 218, 371—373.
- 1868 KNEELAND, SAMUEL, The largest blood discs known. Singular capture of a Canadian Reptile. *Menobranchus lateralis*. *Intellectual Observer*, Vol. 12, 1868, p. 199—200.
- 1896 KNOLL, PH., Ueber die Blutkörperchen bei wechselwarmen Wirbeltieren. *Sitz.-Ber. Kais. Akad. Wiss. Wien*, Bd. 105, 1896, p. 35—66, 3 Taf., 4 Fig.
- 1889 KÖLLIKER, ALBRECHT VON, Handbuch der Gewebelehre des Menschen. Fortgesetzt durch V. VON EBNER. 3 Bde. Leipzig 1889.
- 1903 KÖNIGSTEIN, HANS, Zur Morphologie und Physiologie des Gefäßsystems am Respirationstract. *Anat. Hefte*, Bd. 22, 1903, p. 307—375, 4 Taf., 2 Fig.
- 1903a KÖNIGSTEIN, HANS, Die Funktion der Muskulatur der Amphibienlunge. 1. Anatomischer Teil. *Arch. ges. Physiol.*, Bd. 95, 1903, p. 616—624.
- 1889 KOHL, C., Einige Notizen über das Auge von *Talpa europaea* und *Proteus anguineus*. *Zool. Anz.*, Bd. 12, 1889, p. 383—386, 405—408.
- 1891 KOHL, C., Vorläufige Mitteilung über das Auge von *Proteus anguineus*. *Zool. Anz.*, Bd. 14, 1891, p. 93—96.
- 1864 KORNHUBER, G. A., Bemerkung über das Vorkommen des Olm (*Proteus anguineus* LAUR.). *Verh. d. Ver. f. Naturk. Preßburg*, Bd. 8, 1864, p. 55—57, 1 Fig.
- 1861 KRAINZ, L. M., Weitere Mitteilung über das angebliche Lebendiggebären des *Proteus*. *Verh. d. Zool.-bot. Ges. Wien*, Bd. 12, 1861, p. 37.
- 1879 KUHN, Ueber das häutige Labyrinth der Amphibien. *Arch. mikr. Anat. u. Entw.*, Bd. 17, 1879, p. 479—550.
- 1886 KUHNE, W., Neue Untersuchungen über die motorische Nervenendigung. *Zeitschr. f. Biol.*, Bd. 23, 1886, p. 148, 16 Taf.
- 1903 KUNITOMO, Ueber Blutgefäße in der Epidermis von *Cryptobranchus japonicus*. *Mitt. d. Med. Ges. Tokio*, Bd. 17, 1903.
- 1906 KUPFFER, K. VON, Die Morphogenie des Zentralnervensystems. *Handbuch der Entwicklungslehre der Wirbeltiere* VON OSKAR HERTWIG, Bd. 2, Jena 1906, p. 1—394.
- 1807 LACEPÈDE, M. DE, Sur une espèce de quadrupède ovipare non encore décrite, *Proteus tetradactylus* [*Menobranchus lateralis*]. *Ann. Mus. Hist. nat. Paris*, T. 10, 1807, p. 230—235, 17 pl.
- 1807a LACEPÈDE, M. DE, Sur une espèce de Protée ou Salamandre à quatre doigts à toutes les pattes. *Nouv. Bull. Sci. Soc. Philom.*, T. 1, 1807, p. 64.
- 1906 LANGE, DAN DE, De Kiembladvorming van *Megalobatrachus maximus* SCHL. Amsterdam 1906.
- 1907 LANGE, DAN DE, Die Keimblätterbildung des *Megalobatrachus maximus* (SCHL.). *Anat. Hefte*, Bd. 32, 1907, p. 311—477, 4 Taf.
- 1873 LANGERHANS, PAUL, Notiz zur Anatomie des Amphibienherzens. *Zeitschr. wiss. Zool.*, Bd. 23, 1873, p. 475—458, 1 Fig.
- 1906 LANKES, K., *Necturus maculatus* RAF. in bayerischen Gewässern. *Bl. Aquarien-Kunde*, Magdeburg, Bd. 17, 1906, p. 160—161.
- 1902 LAUBER, H., Anatomische Untersuchung des Auges von *Cryptobranchus japonicus*. *Anat. Hefte*, Bd. 20, 1902, p. 1—17.
- 1826 LE CONTE, JOHN L., Description of a new species of Siren [*striata*] with some observations on animals of a similar nature. *Ann. Lyc. Nat. Hist. New York*, Vol. 1, 1826, p. 52—58.
- 1828 LE CONTE, JOHN L., Description of the Siren *lacertina* n. sp. *Ann. Lyc. Nat. Hist. New York*, Vol. 2, 1828, p. 133—134.
- 1819 LE CONTE, JOHN L., On the habits of *Amphiuma* in a state of captivity. *Proc. Amer. Assoc. Adv. Sci.*, 1819, p. 191—195.
- 1821 LIEBKART, F. S., Einiges über die fischartigen Amphibien. *Isis, Litter. Anz.*, 1821, p. 260, 1 Taf.
- 1810 LIEBKART, F. S., Ueber das Genus *Cryptobranchus*. *FRONIEPS Notizen*, Bd. 13, 1840, p. 19—20.
- 1897 LEVI, GIUS., Ricerche citologiche comparate sulla cellula nervosa dei Vertebrati. *Riv. Pat. nerv. ment. Firenze*, Vol. 2, 1897, p. 193—225, 244—255, 1 tab., 5 fig.

- 1904 LEWIS, FRED T., The question of sinuoids. *Anat. Anz.*, Bd. 25, 1903, p. 261—279, 10 Fig.
- 1853 LEYDIG, F., Anatomisch-histologische Untersuchungen über Fische und Reptilien. Berlin 1853. 4<sup>o</sup>. p. 6 u. 120. 4 Taf.
- 1868 LEYDIG, F., Ueber Organe eines sechsten Sinnes. Zugleich ein Beitrag zur Kenntnis des feineren Baues der Haut bei Amphibien und Fischen. *Nova Acta Acad. Caes. Leopold.* Dresden 1868.
- 1873 LEYDIG, F., Ueber die allgemeinen Bedeckungen der Amphibien. *Arch. mikr. Anat. u. Entw.*, Bd. 9, 1873, p. 753—791, 1 Taf.
- 1876 LEYDIG, F., Ueber die allgemeinen Bedeckungen der Amphibien. *Arch. mikr. Anat. u. Entw.*, Bd. 12, 1876, p. 119—242.
- 1876a LEYDIG, F., Ueber die Schwanzflosse, Tastkörperchen und Endorgane der Nerven bei Batrachiern. *Arch. mikr. Anat. u. Entw.*, Bd. 12, 1876, p. 523.
- 1876b LEYDIG, F., Die Hautdecke und Hautsinnesorgane der Urodelen. *Morphol. Jahrb.*, Bd. 2, 1876, p. 287—318, 4 Taf.
- 1898 LEYDIG, F., Vaskularisiertes Epithel. *Arch. mikr. Anat. u. Entw.*, Bd. 52, 1898, p. 152—155.
- 1766 LINNAEUS, C., *Siren lacertina*. Upsala 1766. 4<sup>o</sup>. p. 15, 1 pl.
- 1866 LLOYD, W. ALFORD, The gigantic Japanese Salamander (*Sieboldia maxima*). *Science Gossip*, 1866, p. 130—131.
- 1905 LOEB, CLARENCE, Some cellular changes in the primary optic vesicles of *Necturus*. *Journ. Comp. Neurol. and Psychol.*, Vol. 15, 1905, p. 459—466, 1 pl.
- 1886 LUCAS, F. A., The sacrum of *Menopoma*. *Amer. Nat.*, Vol. 20, 1886, p. 561—562.
- 1887 MACALLUM, A. B., The termination of nerves in the liver. *Quart. Journ. Micr. Sci.*, Vol. 27, 1887, p. 439—460.
- 1887a MACALLUM, A. B., On the nuclei of the striated muscle-fibre in *Necturus* (*Menobranthus lateralis*). *Quart. Journ. Micr. Sci.*, N. S., Vol. 26, 1887, p. 439—460.
- 1892 MACALLUM, A. B., Studies on the blood of Amphibia. *Trans. Canad. Inst.*, Vol. 2, 1892, Part 2, p. 221—260, 1 pl.
- 1908 Mc GILL, CAROLINE, The fibroglia fibres of *Necturus*. *Anat. Record*, Vol. 2, 1908, p. 146.
- 1908a Mc GILL, CAROLINE, Fibroglia fibres in the intestinal wall of *Necturus* and their relation to myofibris. *Internat. Monatsschr. f. Anat. u. Physiol.*, Bd. 25, 1896, p. 90—98, 1 pl.
- 1896 Mc GREGOR, J. HOWARD, An embryo of *Cryptobranchus*. *Proc. N. Y. Acad. Sci.*, 1896.
- 1896a Mc GREGOR, J. HOWARD, Preliminary notes on the cranial nerves of *Cryptobranchus alleghaniensis*. *Journ. Comp. Neurol. and Psychol.*, Vol. 6, 1896, p. 45—53.
- 1899 Mc GREGOR, J. HOWARD, The spermatogenesis of *Amphiuma*. *Journ. Morphol.*, Vol. 15, 1899, Suppl., p. 57.
- 1875 MALBRANC, M., Bemerkung über die Sinnesorgane der Seitenlinie bei Amphibien. *Centralbl. med. Wiss.*, 1875, No. 1, p. 5.
- 1876 MALBRANC, M., Von der Seitenlinie und ihren Sinnesorganen bei Amphibien. *Zeitschr. wiss. Zool.*, Bd. 26, 1876, p. 24—86.
- 1893 MALL, FRANKLIN P., Histogenesis of the retina in *Amblystoma* and *Necturus*. *Journ. Morphol.*, Vol. 8, 1893, p. 415—432, 12 fig.
- 1839 MANDL, L., Dimensions des globules du sang chez le *Proteus*. *Compt. Rend. Acad. Sc. Paris*, T. 9, 1839, p. 739.
- 1839a MANDL, L., Note sur les globules sanguins du *Protée* et des *Crocodiliens*. *Ann. Sci. nat.*, T. 12, 1839, p. 289—291, fig.
- 1875 MARCHESSETTI, Di alcune nuove località del *Proteus anguineus* LAUR. *Boll. Soc. Adriat. Sc. nat.*, Vol. 1, 1875, p. 192—193.
- 1906 MARCINOWSKI, KATI, Zur Entstehung der Gefäßendothelien und des Blutes bei Amphibien. *Jena. Zeitschr.*, Bd. 41, 1906, p. 19—112, 6 Taf., 17 Fig.
- 1878 MARENZELLER, EMIL VON, Ueber die drei lebenden japanischen Riesensalamander des k. k. Zoologischen Hofkabinetts. *Verh. Zool.-bot. Ges. Wien*, Bd. 27, 1878, p. 43.
- 1893 MARSHALL, WM. B., *Necturus maculatus* in the Hudson River. *Amer. Nat.*, Vol. 26, 1893, p. 779—780.
- 1888 MAURER, F., Die Kiemen und ihre Gefäße bei anuren und urodelen Amphibien. *Morph. Jahrb.*, Bd. 13, 1888, p. 383—384.
- 1888a MAURER, F., Schilddrüse, Thymus und Kiemenreste bei Amphibien. *Morph. Jahrb.*, Bd. 13, 1888, p. 296—382, 3 Taf., 6 Fig.
- 1906 MAURER, F., Die Entwicklung des Muskelsystems und der elektrischen Organe. *Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG*, Bd. 3, Jena 1906.
- 1899 MATHEWS, ALBERT P., The changes in structure of the pancreas cell. *Journ. Morphol.*, Suppl. to Vol. 15, 1899, p. 172—203, 3 pl.
- 1835 MAYER, A. F. J. C., *Analekten für vergleichende Anatomie*. Bonn 1835. 4<sup>o</sup>. p. 6 + 93 + 3, 7 Taf.
- 1819 MECKEL, J. F., Ueber den Darmkanal der Reptilien. *Deutsch. Arch. f. Physiol.*, Bd. 3, 1817, p. 199—232; Nachtrag, Bd. 5, 1819, p. 343—348.
- 1880 MERKEL, FR., Ueber die Endigungen der sensiblen Nerven in der Haut der Wirbeltiere. Rostock 1880.
- 1820 MERREM, BLAS, Versuch eines Systems der Amphibien. *Tentamen Systematis Amphibiorum*. [German and Latin.] Marburg 1820. 8<sup>o</sup>. 1 Taf.

- 1870 METTENHEIMER, C., Einige Betrachtungen über den Olm. Zool. Garten, Bd. 11, 1870, p. 365—368.
- 1893 MEURON, PIERRE DE, Sur la grande Salamandre du Japon. Bull. Soc. Sci. nat. Neuchâtel, T. 21, 1893, p. 186—187.
- 1875 MEYER, FR., Beitrag zur Anatomie des Urogenitalsystems der Selachier und Amphibien. Sitz.-Ber. Naturf. Ges. Leipzig, 1875, p. 38—44.
- 1909 MEYERHOFER, FRANZ, Untersuchungen über die Morphologie und Entwicklungsgeschichte des Rippensystems der urodelen Amphibien. Arb. Zool. Inst. Univ. Wien, 1909, p. 309—352, 2 Taf. u. 9 Fig.
- 1829 MICHAELLES, C., Proteus anguineus Aristoteli prorsus ignotus fuit. Isis, 1829, p. 1270—1273.
- 1831 MICHAELLES, C., Beiträge zur Naturgeschichte des Proteus. Isis, 1831, p. 499—509.
- 1898 MIHALKOVICS, V. VON, Nasenhöhle und JACOBSONSches Organ. Eine morphologische Studie. Anat. Hefte, Bd. 11, 1898, p. 1—107, 11 Taf.
- 1862 MILDE, J., Bemerkungen über den Olm. Verh. d. Zool.-bot. Ges. Wien, Bd. 12, 1862, p. 87—88.
- 1900 MILLER, W. S., The brain of *Necturus maculatus*. Bull. Univ. Wisconsin, 1900, No. 33, p. 227—234, 2 pl.
- 1900a MILLER, W. S., The vascular system of *Necturus maculatus*. Bull. Univ. Wisconsin, Vol. 2, 1900, No. 33, p. 211—226, 2 pl.
- 1900b MILLER, W. S., The lung of *Necturus maculatus*. Bull. Univ. Wisconsin, Vol. 2, 1900, No. 33, p. 203—210, 6 pl.
- 1902 MILLER, W. S., The lymphatics of the lung of *Necturus*. Amer. Journ. Anat., Vol. 2, 1902, p. VI.
- 1905 MILLER, W. S., The blood and lymph vessels of the lung of *Necturus maculatus*. Amer. Journ. Anat., Vol. 4, 1905, p. 445—453, 3 fig.
- 1906 MILLER, W. S., The mesentery in Amphibia and Reptilia. Amer. Journ. Anat., Vol. 4, 1906, p. XIV—XV.
- 1874 MILNER, JAMES W., Report on the Fisheries of the Great Lakes. In United States Commission of Fish and Fisheries. Part 2. Report of the Commissioner for 1872 and 1873. Washington 1874. 8°. p. 78.
- 1892 MINOT, C. S., Human embryology. Philadelphia 1892. 8°.
- 1898 MINOT, C. S., On the veins of the Wolffian body in the Pig. Proc. Boston Soc. Nat. Hist., Vol. 28, 1898, p. 265—274, 1 fig., 1 pl.
- 1900 MINOT, C. S., On a hitherto unrecognized form of blood circulation without capillaries in the organs of Vertebrata. Proc. Boston Soc. Nat. Hist., Vol. 29, 1900, p. 185—215, 12 fig.
- 1822 MITCHILL, SAMUEL L., The Proteus of the North American Lakes. Amer. Journ. Sci. and Arts, Ser. 1, Vol. 4, 1822, p. 181—183.
- 1824 MITCHILL, SAMUEL L., Ueber einige, wahrscheinlich zum Geschlecht der Proteus gehörige Reptilien Nordamerikas. FROBIE'S Not., Bd. 7, 1824, p. 165—166.
- 1824a MITCHILL, SAMUEL L., Observations on several Reptiles of North America which seem to belong to the family Proteus. Amer. Journ. Sci. and Arts, Ser. 1, Vol. 7, 1824, p. 63—69, 2 pl.
- 1869 MIVART, ST. GEORGE, On the myology of *Menopoma alleghaniensis*, *Menobranchnus lateralis* and *Chamaeleon Parsonii*. Proc. Zool. Soc. London, 1869 and 1870.
- 1869a MIVART, ST. GEORGE, Notes on the myology of *Menobranchnus lateralis*. Proc. Zool. Soc. London, 1869.
- 1870 MIVART, ST. GEORGE, On the axial skeleton of the Urodela. Proc. Zool. Soc. London, 1870.
- 1887 MOESCH, C., Der japanische Riesensalamander (*Cryptobranchnus japonicus*) und der fossile Salamander von Oeningen (*Andrias scheuchzeri*). Neujahrsblatt Naturf. Ges. Zürich, 1887. 4°. p. 12.
- 1880 MONTGOMERY, HENRY, Some observations on the *Menobranchnus maculatus*. Canadian Naturalist, Vol. 9, 1880, p. 160—164.
- 1898 MONTE, RINA, Su la morfologia comparata dei condotti escretori delle ghiandole gastriche nei Vertebrati. Boll. Sc. Pavia, 1898, Anno 20, p. 33—39, 65—75, 101—108, 2 tab.
- 1899 MONTE, RINA, Ricerche anatomo-comparative sulla minuta innervazione degli organi trofici nei Cranioti inferiori. Torino 1899. 147 pp., 12 pl.
- 1908 MOODIE, ROY L., The ancestry of the caudate Amphibia. Amer. Nat., Vol. 42, 1908, p. 361—373.
- 1897 MORGAN, T. H., The development of the Frog's egg. New York 1897. 8°.
- 1903 MORGAN, T. H., Regeneration of the leg of *Amphiuma means*. Biol. Bull., Vol. 5, 1903, p. 293—296.
- 1832 MÜLLER, JOH. CHRISTOF, Beiträge zur Anatomie und Naturgeschichte der Amphibien. Zeitschr. Physiol., Bd. 4, 1832.
- 1864 MÜLLER, KARL, Der Riesensalamander. Natur, Bd. 13, 1864, p. 172—174.
- 1897 MURRAY, J. A., The vertebral column of certain primitive Urodela. Anat. Anz., Bd. 13, 1897, p. 661—664, 3 Fig.
- 1891 NAKAGAWA, ISAAC, The origin of the cerebral cortex and homologies of the optic lobe layers in the lower vertebrates. Journ. Morphol., Vol. 4, 1891, p. 1—10, 1 pl.
- 1828 NEILL, PATRICK, Some account of the habits of a specimen of *Siren lacertina* kept alive at Canonmills. Edinb. New Phil. Journ., Vol. 4, 1828, p. 346—355.
- 1832 NEILL, PATRICK, Siren. Isis, 1832, p. 698—699.
- 1902 NEMILOFF, A., Zur Frage über die Nerven des Darmkanales bei den Amphibien. Schriften Kais. Naturf. Ges. St. Petersburg, Bd. 32, 1902, p. 59—96.

- 1906 NEUMAYER, L., Histogenese und Morphogenese des peripheren Nervensystems der Spinalganglien und des Nervus sympathicus. *Handbuch der Entwicklungslehre der Wirbeltiere* von OSKAR HERTWIG, Bd. 2, Jena 1906, p. 573—621.
- 1893 NICLOGU, PHILIPPOS, Ueber die Hautdrüsen der Amphibien. *Zeitschr. wiss. Zool.*, Bd. 56, 1893, p. 109—187, 3 Taf.
- 1864 NOBBE, F., *Proteus anguineus* in der Gefangenschaft. *Zool. Garten*, Bd. 5, 1864, p. 231.
- 1894 NORRIS, H. W., The development of the auditory vesicle in *Necturus*. *Proc. Iowa Acad. Sci.*, Vol. 1, 1894, p. 105—107.
- 1905 NORRIS, H. W., The so-called Dorso-trachealis branch of the seventh cranial nerve in *Amphiuma*. *Anat. Anz.*, Bd. 27, 1905, p. 271—272.
- 1908 NORRIS, H. W., The cranial nerve components in *Amphiuma*. *Science*, N. S. Vol. 27, 1908.
- 1909 NORRIS, H. W., The cranial nerves of *Amphiuma means*. *Journ. Comp. Neurol. and Psychol.*, Vol. 18, 1909, p. 527—568, 5 pl.
- 1907 NUSBAUM, J., Ein Fall von Viviparität beim *Proteus anguineus*. *Biol. Centralbl.*, Bd. 27, 1907, p. 370—375.
- 1908 NUSBAUM, J., Zur Entwicklungsgeschichte der Occipitalregion des Schädels und der WEBERSchen Knochelehen bei den Cyprinoiden. Krakau (Acad.) 1908. 8°.
- 1789 OESTERDAM, ABRAHAM, *Siren lacertina*. *Amoenitates Academicae*, Vol. 7, 1789, p. 322.
- 1906 OKAJIMA, KEYI, Zur Anatomie des inneren Gehörganges von *Cryptobranchus japonicus*. *Anat. Hefte*, Bd. 32, 1906, p. 233.
- 1906 OKAJIMA, KEYI, Zur Anatomie des Geruchsorgans von *Cryptobranchus japonicus*. *Anat. Anz.*, Bd. 29, 1906, p. 641—650, 5 Fig.
- 1817 OKEN, LORENZ, Ueber den Olm. *Proteus anguineus*. *Isis*, 1817, p. 641—645, 1 Fig.
- 1836 OKEN, LORENZ, *Allgemeine Naturgeschichte*, Bd. 6, Stuttgart 1836, p. 417. 8°.
- 1890 OPPEL, A., Beiträge zur Anatomie des *Proteus anguineus*. *Arch. mikr. Anat. u. Entw.*, Bd. 34, 1890.
- 1895 OPPEL, A., Die Magendrüsen der Wirbeltiere. *Anat. Anz.*, Bd. 11, 1895, p. 596—601.
- 1896 OPPEL, A., *Lehrbuch der vergleichenden mikroskopischen Anatomie der Wirbeltiere*. Jena 1896.
- 1899 OSAWA, G., Beiträge zur Anatomie des japanischen Riesensalamanders. *Festschrift TAGUCHI*, Tokio 1899.
- 1902 OSAWA, G., Beiträge zur Anatomie des japanischen Riesensalamanders. *Mitt. Med. Facult. Tokio*, Vol. 5, 1902, p. 8—40, 7 Taf.
- 1883 OSBORN, H. F., Preliminary observations upon the brain of *Amphiuma*. *Proc. Acad. Nat. Sci. Philadelphia*, 1883, p. 177—184.
- 1884 OSBORN, H. F., Preliminary notes upon the brain of *Menopoma*. *Proc. Acad. Nat. Sci. Philadelphia*, 1884.
- 1886 OSBORN, H. F., Preparing brain of Urodela. *Journ. Roy. Micr. Soc.*, (2) Vol. 5, 1886, p. 536—537.
- 1886a OSBORN, H. F., The origin of the corpus callosum; a contribution upon the cerebral commissures of the vertebrata. *Morphol. Jahrb.*, Bd. 12, 1886, p. 223—251.
- 1888 OSBORN, H. F., A contribution to the internal structure of the Amphibian brain. *Journ. Morph.*, Vol. 2, 1888, p. 51—92, 3 pl.
- 1884 OSBORN, J., Observations upon the urodele Amphibian brain. *Zool. Anz.*, Bd. 7, 1884, p. 679—682.
- 1835 OWEN, RICHARD, On the structure of the heart in the Perennibrancheata. *Trans. Zool. Soc. London*, 1835.
- 1842 OWEN, RICHARD, On the blood-disks in *Siren lacertina*. *Micr. Journ.*, 1842.
- 1866 OWEN, RICHARD, On the anatomy of Vertebrates. Vol. 3. London 1866. 8°.
- 1896 PARKER, G. H., Variations in the vertebral column of *Necturus*. *Anat. Anz.*, Bd. 11, 1896, p. 711—717, 2 Fig.
- 1868 PARKER, W. K., A monograph on the structure and development of the shoulder girdle and sternum in the Vertebrata. Ray Soc. London, 1868.
- 1877 PARKER, W. K., On the structure and development of the skull in the Urodela Amphibia. *Phil. Trans. Roy. Soc. London*, Vol. 167, 1877, p. 529—597, 9 pl.
- 1894 PETER, KARL, Ueber die Bedeutung des Atlas der Amphibien. *Anat. Anz.*, Bd. 10, 1894, p. 565—574.
- 1898 PETER, KARL, Die Entwicklung und funktionelle Gestaltung des Schädels von *Ichthyophis glutinosus*. *Morphol. Jahrb.*, Bd. 25, 1898, p. 555—628, 3 Taf., 1 Fig.
- 1897 PHISALIX, C., Action physiologique du venin de Salamandre du Japon (*Sieboldia maxima*). Atténuation par la chaleur et vaccination de la grenouille contre ce venin. *Compt. Rend. Soc. Biol. Paris*, T. 4, 1897, p. 723—725.
- 1897a PHISALIX, C., Propriétés immunisantes du venin de Salamandre du Japon vis-à-vis du venin de vipère. *Compt. Rend. Soc. Biol. Paris*, T. 4, 1897, p. 822—823.
- 1889 PIERSOL, G. A., Structure of spermatozoa [*Amphiuma tridactylum*]. *Univ. Med. Magaz. Philad.*, 1889, p. 20.
- 1894 PLATT, JULIA B., Ontogenetische Differenzierung des Ektoderms in *Necturus*. 1. Studie. *Arch. f. mikr. Anat. u. Entw.*, Bd. 43, 1894, p. 911—966, 6 Taf.
- 1896 PLATT, JULIA B., The development of the thyroid gland and of the suprapericardial bodies in *Necturus*. *Anat. Anz.*, Bd. 11, 1896, p. 557—567, 9 Fig.

- 1896a PLATT, JULIA B., Ontogenetic differentiations of the ectoderm in *Necturus*. Study 2. On the development of the peripheral nervous system. *Quart. Journ. Micr. Sci.*, Vol. 38, 1896, p. 485—547, 2 pl.
- 1897 PLATT, JULIA B., Development of the cartilaginous skull and of the branchial and hypoglossal musculature in *Necturus*. *Morphol. Jahrb.*, Bd. 35, 1897, p. 377—464, 3 pl.
- 1859 POMPE VAN MEERDERVOORT, J. L. C., Over den grooten Japanschen Salamander. *Natuurk. Tijdschr. Nederl. Indië*, Ser. 4, Vol. 6, 1859, p. 386.
- 1904 PRENANT, A., Sur la morphologie des cellules épithéliales ciliées qui recouvrent le péritoine hépatique des Amphibiens. *Compt. Rend. Soc. Biol. Paris*, T. 55, 1903, p. 1044—1046.
- 1895 RABL, H., Ueber die Herkunft des Pigments in der Haut der Larven der urodelen Amphibien. *Anat. Anz.*, Bd. 10, 1895, p. 12—16.
- 1889 RABL, CARL, Theorie des Mesoderms. *Morphol. Jahrb.*, Bd. 15, 1889, p. 113—252, 4 Taf., 9 Fig.
- 1898 RABL, CARL, Ueber den Bau und die Entwicklung der Linse. 1. Selachier und Amphibien. *Zeitschr. wiss. Zool.*, Bd. 63, 1898, p. 486—572, 4 Taf., 14 Fig.
- 1819 RAFINESQUE, Prodrôme de soixante-dix nouveaux genres d'animaux découverts dans l'intérieur des Etats Unis d'Amérique durant l'année 1818. *Journ. de Phys.*, T. 88, 1819, p. 417—429.
- 1825 RATHKE, H., Ueber die Entwicklung der Geschlechtssteile bei den Amphibien. *Beiträge zur Geschichte der Tierwelt*. 3 Abt. Halle 1825.
- 1904 REESE, A. M., The sexual elements of the Giant Salamander. *Amer. Nat.*, Vol. 38, 1904, p. 497.
- 1904a REESE, A. M., The sexual elements of the Giant Salamander, *Cryptobranchus alleghaniensis*. *Biol. Bull.*, Vol. 6, 1904, p. 220—223.
- 1904b REESE, A. M., The enteron and integument of *Cryptobranchus alleghaniensis*. *Trans. Amer. Micr. Soc.*, Vol. 26, 1904, p. 109—120.
- 1905 REESE, A. M., Integument von *Cryptobranchus*. *Trans. Amer. Micr. Soc.*, Vol. 26, 1905, p. 109—120, 2 pl.
- 1905a REESE, A. M., The eye of *Cryptobranchus*. *Biol. Bull.*, Vol. 9, 1905, p. 22—26, 1 fig.
- 1906 REESE, A. M., Anatomy of *Cryptobranchus alleghaniensis*. *Amer. Nat.*, Vol. 40, 1906, p. 287—326.
- 1906a REESE, A. M., Observations on the reactions of *Cryptobranchus* and *Necturus* to light and heat. *Biol. Bull.*, Vol. 11, 1906, p. 93—99.
- 1876 REIN, J. J., and RORETZ, A. VON, Beitrag zur Kenntnis des Riesensalamanders (*Cryptobranchus japonicus*). *Zool. Garten*, Bd. 17, 1876, p. 33—37.
- 1884 REIN, J. J., Reptiles and batrachians [Japan]. New York 1884. p. 189.
- 1897 REJSEK, J., Histologie oka od *Cryptobranchus japonicus*. *Rozpr. Ceské. Akad.*, Ser. 2, 1897, p. 10.
- 1897a REJSEK, J., L'histologie de l'oeil de *Cryptobranchus japonicus*. *Bibliogr. Anat.*, Année 5, 1897, p. 139—146.
- 1881 RETZIUS, G., Das Gehörorgan der Wirbeltiere. 1. Das Gehörorgan der Fische und Amphibien. Stockholm 1881.
- 1904 RETZIUS, G., Zur Kenntnis der Limitans externa der nervösen Zentralorgane. *Biol. Unters. RETZIUS*, Bd. 11, 1904, p. 77—81.
- 1905 RETZIUS, G., Zur Kenntnis der Nervenendigungen in den Papillen der Zunge der Amphibien. *Biol. Unters. RETZIUS*, Bd. 12, 1905, p. 61—64.
- 1897 REYNOLDS, SIDNEY H., The vertebrate skeleton. Cambridge 1897. 8<sup>o</sup>.
- 1907 RIBBING, L., Die distale Armmuskulatur der Amphibien, Reptilien und Säugetiere. *Zool. Jahrb.*, Bd. 23, 1907, p. 587—683, 2 Taf.
- 1902 RICHTER, ALEX V., Ueber die Struktur und die Bedeutung der LANGERHANS'schen Inseln im Pancreas der Amphibien. Berlin 1902. p. 37.
- 1909 RIDDLE, OSCAR, The rate of digestion in cold-blooded vertebrates. The influence of season and temperature. *Amer. Journ. Physiol.*, Vol. 24, 1909, p. 447—458.
- 1905 ROMEISER, THEODORE H., A case of abnormal venous system in *Necturus maculatus*. *Amer. Nat.*, Vol. 39, 1905, p. 391—396.
- 1907 ROOY, PETRONELLA JOHANNA DE, Die Entwicklung des Herzens, des Blutes und der großen Gefäße bei *Megalobatrachus maximus* SCHLEGEL. *Jena. Zeitschr.*, Bd. 35, 1907, p. 309—346, 6 Taf.
- 1901 RUBASCHKIN, W. J., Ueber den Bau des Gehirns der Amphibien. XI. Vers. russ. Naturf. u. Aerzte, St. Petersburg 1901.
- 1903 RUBASCHKIN, W. J., Zur Morphologie des Gehirns der Amphibien. *Arch. mikr. Anat. u. Entw.*, Bd. 62, 1903, p. 207—213, 2 Taf.
- 1817 RUDOLPH, K. A., Ueber den *Proteus anguineus*. *Isis*, 1817, p. 1017—1019.
- 1897 RUGE, GEORG, Ueber das peripherische Gebiet des Nervus facialis bei Wirbeltieren. *Festschrift GEGENBAUR*, Bd. 3, 1897, p. 193—348, 76 Fig.
- 1818 RUSCONI, M., and CONFIGLIACHI, Del *Proteo anguineo* di LAURENTI monografia. Pavia 1818.
- 1827 RUSCONI, MAURO, Descrizione di un *Proteo femmina* notabile per lo sviluppo delle parti della generazione. *Isis*, 1827, p. 94—100.
- 1828 RUSCONI, MAURO, Sopra un *Proteo femineo*. Pavia 1828. 4<sup>o</sup>. 1 tab.

- 1837 RUSCONI, MAURO, Observations anatomiques sur la Sirène mise en parallèle avec le Protée et le têtard de la Salamandre aquatique. Pavie 1837. 4<sup>o</sup>. 6 pl.
- 1843 RUSCONI, MAURO, Osservazioni sopra il Proteo. Milano 1843. 8<sup>o</sup>.
- 1843a RUSCONI, MAURO, Nuove osservazione sopra il Proteo anguineo di LAURENTI. Giorn. dell'Istit. Lomb. e Bibl. Ital., Vol. 6, 1843, p. 288—290.
- 1879 RYDER, JOHN A., Morphological notes on the limbs of the Amphiumidae, as indicating a possible synonymy of the supposed genera. Proc. Acad. Sci. Philad., 1879, p. 14—15.
- 1890 RYDER, JOHN A., On a brood of larval Amphiuma. Amer. Nat., Vol. 23, 1890, p. 927—928.
- 1903 SAINT-HILAIRE, CONST., Bau des Darmepithels bei Amphiuma. Anat. Anz., Bd. 22, 1903, p. 489—493, 6 Fig.
- 1887 SASAKI, C., Some notes on the Giant Salamander of Japan. Journ. Coll. of Sci. Tokyo, Japan, Vol. 1, 1887, p. 269—274.
- 1902 SCHAUER, A., Ueber kontraktile Fibrillen in den glatten Muskelfasern des Mesenteriums der Urodelen. Anat. Anz., Bd. 22, 1902, p. 65—82, 2 Taf., 6 Fig.
- 1906 SCHAUNSLAND, H., Die Entwicklung der Wirbelsäule, nebst Rippen und Brustbein. Handbuch der Entwicklungslehre der Wirbeltiere von OSKAR HERTWIG, Bd. 3, 1906, p. 339—572.
- 1891 SCHLAMPP, K. W., Die Augenlinse des Proteus anguineus. Biol. Centralbl., Bd. 11, 1891, p. 40—42.
- 1892 SCHLAMPP, K. W., Das Auge des Grottenolmes (*Proteus anguineus*). Zeitschr. wiss. Zool., Bd. 53, 1892, p. 537—557, 1 Taf.
- 1837 SCHLEGEL, H., Abbildungen neuer oder unvollständig bekannter Amphibien nach der Natur oder dem Leben entworfen. Düsseldorf 1837—1844. kl. Fol. 50 Taf.
- 1838 SCHLEGEL, H., Fauna Japonica. Lugd. Bat. 1838, p. 127, 2 tab.
- 1850 SCHMIDL, A., Notizen über die von ihm aus der Plauina-Höhle mitgebrachten und der Klasse vorgezeigten Proteen. Sitz.-Ber. math.-nat. Akad. Wiss. in Wien, Bd. 5, p. 228—232.
- 1864 SCHMIDT, F. J. J., GODDARD, Q. J., en VAN DER HOEVEN, J., Aanteekeningen over de anatomie van den *Cryptobranchus japonicus*. Natuurk. Verh. Maatsch. Wetensch. Haarlem, Vol. 19, 1864, p. 66, 12 pl.
- 1878 SCHMIDT, H. D., The structure of the coloured bloodcorpuscles of *Amphiuma tridactylum*, the frog and man. Journ. Roy. Micr. Soc., Vol. 1, 1878, p. 57—78 and 97—120.
- 1904 SCHMIDT, VICTOR, Studien über Ovogenese. 1. Die Wachstumsperiode der Eier von *Proteus anguineus*. Anat. Hefte, Bd. 27, 1904, p. 1—71, 4 Taf.
- 1779 SCHNEIDER, J. G., Historiae Amphibiorum naturales et litterariae. Fasciculus primus. Jenae 1779. 8<sup>o</sup>.
- 1873 SCHNEIDER, O., Ueber den Olm. Gaea, Bd. 9, 1873, p. 524.
- 1890 SCHNEIDER, R., Neue histologische Untersuchungen über die Eisenaufnahme in den Körper des Proteus. Sitz.-Ber. K. Preuß. Akad. Wiss. Berlin, Bd. 36, 1890, p. 887—897.
- 1904 SCHOENICHEN, WALTHER, Die Brutpflege bei den Amphibien und besonders bei dem japanischen Riesensalamander (*Megalobatrachus maximus*). Prometheus, Bd. 16, 1904, p. 37—40, 52—54, 15 Fig.
- 1801 SCHREIBERS, CH., A historical and anatomical description of a doubtful amphibious animal of Germany called by LAURENTI *Proteus anguineus*. Phil. Trans. Roy. Soc. London, Vol. 91, 1801, p. 24, 2 pl.
- 1818 SCHREIBERS, CH., *Proteus anguineus*. Viennae 1818. 4<sup>o</sup>.
- 1820 SCHREIBERS, CH., Sur le Protée. Isis, 1820, p. 567—570.
- 1893 SCHUBERG, A., Beiträge zur Kenntnis der Amphibienhaut. Zool. Jahrb., Bd. 6, 1893, p. 481—490, 1 Taf.
- 1900 SCHULZ, PAUL, Ueber die Anordnung der Muskulatur im Magen der Batrachier. Arch. Anat. Physiol., Phys. Abt., 1900, p. 1—8, 4 Fig.
- 1861 SCHULZE, F. E., Ueber die Nervenendigung in den sogenannten Schleimkanälen der Fische und über entsprechende Organe der durch Kiemen atmenden Amphibien. Arch. Anat. Physiol., 1861, p. 759—769, 1 Taf.
- 1870 SCHULZE, F. E., Ueber die Sinnesorgane der Seitenlinie bei Fischen und Amphibien. Arch. mikr. Anat. u. Entw., Bd. 6, 1870, p. 62, 3 Taf.
- 1876 SCHULZE, F. E., Zur Fortpflanzungsgeschichte des *Proteus anguineus*. Zeitschr. wiss. Zool., Bd. 26, 1876, p. 350—354.
- 1905 SCHULZE, O., Ueber partiell albinotische und mikrophthalmische Larven von *Salamandra maculata* nebst einigen Angaben über die Fortpflanzung dieses Tieres. Zeitschr. wiss. Zool., Bd. 82, 1905, p. 472—493.
- 1905 SEYDEL, O., Ueber die Nasenhöhle und das JACOBSONSCHE Organ der Amphibien. Morphol. Jahrb., Bd. 23, 1895, p. 453—543, 22 Fig.
- 1828 SIEBOLD, CAROL TH. ERNST, Observationes quaedam de salamandris et tritonibus. Berolini 1828. p. 6 + 30, 1 tab.
- 1833 SIEBOLD, PHILIPP FRANZ VON, Fauna japonica. LEYDEN 1833—1851.
- 1904 SIEGLBAUER, F., Zur Anatomie der Urodelenextremität. Arch. f. Anat. u. Entw., 1904, p. 385—404, 1 Taf.
- 1908 SMALLWOOD, W. M., The sacrum of *Necturus*. Anat. Anz., Bd. 33, 1908, p. 237—239, 1 Fig.
- 1890 SMIRNOW, A., Die Struktur der Nervenzellen im Sympathicus der Amphibien. Arch. mikr. Anat. u. Entw., Bd. 35, 1890, p. 407—424, 2 Taf.
- 1900 SMITH, F., Some additional data on the position of the sacrum in *Necturus*. Amer. Nat., Vol. 34, 1900, p. 635—638.

- 1906 SMITH, BERTRAM, G., Preliminary report on the embryology of *Cryptobranchus alleghaniensis*. Biol. Bull., Vol. 9, 1906, p. 146—164, 1 pl.
- 1898 SMITH, HUGH M., On the occurrence of *Amphiuma*, the so-called Congo Snake in Virginia. Proc. U. S. Nat. Mus., Vol. 21, 1898, p. 379—380.
- 1832 SMITH, J., Isis, 1832, p. 1088.
- 1828 SMITH, F. AUG., Account of the dissection of a *Proteus* of the Lakes (*Menobranchus*) with remarks on the *Siren intermedia*. Ann. Lye. Nat. Hist. New York, Vol. 2, 1828, p. 259—263.
- 1897 SOBOTTA, J., Die Furchung des Wirbeltiereies. Ergeb. Anat. Entw., Bd. 6, 1897, p. 493—593, 38 Fig.
- 1902 SOBOTTA, J., Ueber die Entwicklung des Blutes, des Herzens und der großen Gefäßstämme der Salmoniden, nebst Mitteilungen über die Ausbildung der Herzform. Anat. Hefte, 1902, Heft 63, p. 579—688.
- 1869 SPICER, W. W., *Proteus anguineus*. Sci. Gossip, 1869, p. 135.
- 1892 STEINER, LEONHARD, Preliminary description of a new genus and species of blind cave Salamander from North America (*Typhlotriton spelaeus*). Proc. U. S. Nat. Mus., Vol. 15, 1892, p. 115—117, 1 pl.
- 1896 STEINER, LEONHARD, Description of a new genus and species of blind tailed Batrachians from the subterranean waters of Texas. Proc. U. S. Nat. Mus., Vol. 18, 1896, p. 619—621.
- 1907 STEINER, LEONHARD, Herpetology of Japan and adjacent territory. Washington 1907. 8°.
- 1907a STEINER, LEONHARD, Specific name of *Necturus maculosus*. Science, Vol. 25, 1907, p. 190.
- 1906 STOHK, PHILIPP, Histology arranged upon an embryological basis by FREDERIC T. LEWIS. Philadelphia 1906.
- 1870 STRAUCH, ALEXANDER, Revision der Salamandriden-Gattungen nebst Beschreibung einiger neuen oder weniger bekannten Arten dieser Familie. Mem. Ac. Sc. St. Pétersburg, T. 16, 1870, p. 109, 2 pl.
- 1872 STRICKER, S. (ed.), Manual of human and comparative histology. Tr. by HENRY POWER. 3 vols. London 1872.
- 1892 STRONG, OLIVER S., The structure and homologies of the cranial nerves of the Amphibia as determined by their peripheral distribution and internal origin. Anat. Anz., Bd. 7, 1892, p. 467—471.
- 1895 STRONG, OLIVER S., The cranial nerves of Amphibia. Journ. Morphol., Vol. 10, 1895, p. 101—230, 6 pl.
- 1888 STRUBE, K. W., [Ueber den japanischen Salamander]. [Russian.] Nachr. d. Kais. Ges. d. Freunde d. Naturwiss. Moskau, Bd. 50, 1888.
- 1874 STUDER, TH., Ueber die Epidermis der *Amphiuma didactyla* [means]. Mitt. Naturf. Ges. Bern, 1874, p. 48.
- 1897 STUDNÍČKA, F. K., Studien über den Bau des Sehnerven der Wirbeltiere. Jen. Zeitschr., Bd. 31, 1897, p. 1—28, 2 Taf.
- 1909 TAKAHASHI, KATASHI, Histogenesis of the lateral line system in *Necturus*. Thesis, Chicago 1909.
- 1838 TEMMINCK, C. J., and SCHLEGEL, H., Fauna Japonica auctore PH. FR. DE SIEBOLD. Reptilia elaborantibus C. J. TEMMINCK et H. SCHLEGEL. Cum mappa geographico-zoologica et tabulis lithogr. XXVIII. Lugduni Batavorum, ex officin. lithogr. auctoris et typis J. G. Lalau, 1838. Fol. XXII + 144 pp. + doublepage map + 9 + 10 + 8 pl.
- 1904 TERRY, ROBERT J., Two skulls of larval *Necturus*. Amer. Journ. Anat., Vol. 3, 1904, p. XI.
- 1906 TERRY, ROBERT J., The nasal skeleton of *Amblystoma punctatum* (LIN.). Trans. St. Louis Acad. Sci., Vol. 16, 1906, p. 95—124, 4 pl.
- 1875 TOMMASINI, M., Sulla diffusione del *Proteus*. Bollet. Soc. Adriat. Sc. nat., Vol. 1, 1875, p. 152—156.
- 1882 TOWNSEND, CHAS. H., Habits of *Menopoma*. Amer. Nat., Vol. 16, 1882, p. 139—140.
- 1820 TREVIANUS, De protei anguinei encephalo et organis sensuum disquisitiones zootomicae. Commentationes societatis regiae scientiarum Göttingensis recentiores, Göttingen 1820, Vol. 4.
- 1837 TSCHUDI, J. J., Ueber den Homo diluvii testis (*Andrias Scheuchzeri*). Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde, Stuttgart 1837, Heft 5, p. 545—547.
- 1838 TSCHUDI, J. J., Classification der Batrachier mit Berücksichtigung der fossilen Tiere dieser Abteilung der Reptilien. Mem. Soc. Sci. nat. de Neuchâtel, Vol. 2, 1838.
- 1903 TUEBERCKHEIM, W., Ueber das Rückenmark von *Cryptobranchus japonicus*. Leipzig 1903. p. 21, 9 Taf.
- 1862 VAILLANT, LÉON, Note sur la structure du noyau des globules sanguins, et la composition de l'encéphale chez la Sirène lacertine. Compt. Rend. et Mém. Soc. d. Biol. Paris, Sér. 3, T. 4, 1862, p. 4—6.
- 1863 VAILLANT, LÉON, Mémoire pour servir à l'histoire anatomique de la sirène lacertine. Ann. Sci. nat. Zool., Sér. 4, T. 19, 1863, p. 295—316, 3 pl.
- 1863a VAILLANT, LÉON, Note sur l'anatomie de la sirène lacertine. Compt. Rend. Acad. Sc. Paris, T. 56, 1863, p. 839.
- 1837 VALENTIN, G., Bruchstücke aus der feineren Anatomie des *Proteus anguineus*. Repert. f. Anat. u. Physiol., Bd. 1, 1837, p. 282—291.
- 1841 VALENTIN, G., Ueber die Samentierbündel und die Afterdrüse des *Proteus anguineus*. Repert. f. Anat. u. Physiol., Bd. 6, 1841, p. 353—358.
- 1903 VAN PEE, P., Recherches sur le développement des extrémités chez *Amphiuma* et *Necturus*. Compt. Rend. Assoc. Anat. Sess. 5. Liège, 1903, p. 37—42.
- 1903a VAN PEE, P., Ueber die Entwicklung der Extremitäten bei *Amphiuma* und *Necturus*. Verh. Anat. Ges. Heidelberg, 1903, p. 83—86.

- 1904 VAN PEE, P., Les membres chez *Amphiuma*. *Anat. Anz.*, Bd. 24, 1904, p. 176—182, 1 Fig.
- 1837 VIATOR (?), On the *Proteus anguineus*. *CHARLESWORTH'S Magaz. Nat. Hist.*, Vol. 1, 1837, p. 625—630.
- 1894 VIRCHOW, HANS, Einige embryologische und angiologische Erfahrungen über nordamerikanische Wirbeltiere. *Sitz.-Ber. Ges. naturf. Freunde Berlin*, 1894, p. 33—41.
- 1904 VIRÉ, A., Sur quelques expériences effectuées au laboratoire des Catacombes du Muséum d'Histoire naturelle. *Compt. Rend. Acad. Sci. Paris*, T. 138, 1904, p. 706—708.
- 1828 WAGLER, JOH., *Descriptiones et icones Amphibiorum*. Stuttgartiae 1828—1833. fol. 36 tab.
- 1830 WAGLER, JOH., *Natürliches System der Amphibien mit vorgehender Classification der Säugetiere und Vögel*. München 1830. 8<sup>o</sup>.
- 1880 WAGNER, FRANZ, Der Olm (*Proteus anguineus*). *Isis Russ.*, Bd. 5, 1880, p. 411—412.
- 1837 WAGNER, RUD., Notes on *Proteus anguineus*. *Proc. Zool. Soc. London*, Vol. 5, 1837, p. 107—108.
- 1897 WAITE, F. C., Variations in the brachial and lumbo-sacral plexi of *Necturus maculosus* RAFINESQUE. *Bull. Mus. Comp. Zool. Harvard Univ.*, Vol. 31, 1897, p. 69—92.
- 1907 WAITE, F. C., Specific name of *Necturus maculosus*. *Amer. Nat.*, Vol. 41, 1907, p. 23—30.
- 1887 WALTER, FERDINAND, Das Visceralskelett und seine Muskulatur bei den einheimischen Amphibien und Reptilien. *Jen. Zeitschr.*, Bd. 21, 1887, p. 1—45, 4 Taf.
- 1905 WARREN, JOHN, The development of the paraphysis and the pineal region in *Necturus maculatus*. *Amer. Journ. Anat.*, Vol. 5, 1905, p. 1—27, 23 fig.
- 1881 WEINZETTL, V., Zur Kenntniss des weiblichen *Proteus anguineus*. *Sitz.-Ber. K. böhm. Ges. Prag*, 1881, p. 297—303, 1 Taf.
- 1878 WEISMANN, A., and WIEDERSHEIM, R., Zwei noch junge *Siren lacertina*. Aus dem zoologischen und anatomischen Institut der Universität Freiburg i. Br. *Zool. Anz.*, Bd. 1, 1878, p. 6.
- 1890 WELIKI, W. N., Ueber Mehrzahl der Lymphherzen bei *Proteus anguineus* und *Rana temporaria*. *Trav. Soc. Natur. St. Pétersbourg, Sect. Zool.*, Vol. 20, 1890.
- 1906 WHIPPLE, INEZ, The ypsiloid apparatus of Urodeles. *Biol. Bull.*, Vol. 10, 1906, p. 255—297.
- 1885 WHITMAN, C. O., *Methods of research in microscopical anatomy and embryology*. Boston 1885. 8<sup>o</sup>. p. 255.
- 1888 WHITMAN, C. O., The eggs of Amphibia. *Amer. Nat.*, Vol. 22, 1888, p. 857.
- 1888a WHITMAN, C. O., Some new facts about the Hirudinea. *Journ. Morphol.*, Vol. 2, 1888, p. 585—599.
- 1899 WHITMAN, C. O., *Animal behavior. Biological Lectures from the Marine Biological Laboratory Woods Holl, Mass.*, 1898, Boston. 8<sup>o</sup>. p. 285—338.
- 1896 WHITMAN, C. O., and EYCLESHYMER, A. C., The egg of *Amia* and its cleavage. *Journ. Morphol.*, Vol. 12, 1896, p. 309—356, 1 pl.
- 1865 WIED-NEUWIED, PRINZ MAXIMILIAN ZU, Verzeichnis der Reptilien, welche auf einer Reise im nördlichen Amerika beobachtet wurden. *Nov. Acta Acad. Leop.-Carol.*, Vol. 32, 1865.
- 1877 WIEDERSHEIM, R., Zur Fortpflanzungsgeschichte des *Proteus anguineus*. *Morphol. Jahrb.*, Bd. 3, 1877, p. 632.
- 1877a WIEDERSHEIM, R., Das Kopfskelett der Urodelen. *Morphol. Jahrb.*, Bd. 3, 1877, p. 352—448, 5 Taf., 1 Fig.: p. 459—548, 4 Taf., 5 Fig.
- 1877b WIEDERSHEIM, R., Ueber Neubildung von Kiemen bei *Siren lacertina*. *Morphol. Jahrb.*, Bd. 3, 1877, p. 630—632.
- 1890 WIEDERSHEIM, R., Beiträge zur Entwicklungsgeschichte von *Proteus anguineus*. *Arch. mikr. Anat. u. Entw.*, Bd. 35, 1890, p. 121—140, 2 Taf.
- 1892 WIEDERSHEIM, R., Das Gliedmaßenskelett der Wirbeltiere mit besonderer Berücksichtigung des Schulter- und Beckengürtels bei Fischen, Amphibien und Reptilien. Mit 40 Figuren im Texte und einem Atlas von 17 Tafeln. Jena 1892. 8<sup>o</sup>. p. 267.
- 1909 WIEDERSHEIM, R., *Lehrbuch der vergleichenden Anatomie der Wirbeltiere*. 1909.
- 1874 WILDER, B. G., *Menobranchus edible*. *Amer. Nat.*, Vol. 8, 1874, p. 438.
- 1883 WILDER, B. G., On the habits of *Cryptobranchus*. *Amer. Nat.*, Vol. 16, 1883, p. 816—817.
- 1891 WILDER, H. H., A contribution to the anatomy of *Siren lacertina*. *Zool. Jahrb.*, Bd. 4, 1891, p. 653—691.
- 1892 WILDER, H. H., Die Nasengegend von *Menopoma alleghaniense* und *Amphiuma tridactylum* nebst Bemerkungen über die Morphologie des Ramus ophthalmicus profundus trigemini. *Zool. Jahrb., Abt. I. Anat. u. Ontog.*, Bd. 5, 1892, p. 155—176, 2 Taf.
- 1896 WILDER, H. H., The amphibian larynx. *Zool. Jahrb.*, Bd. 9, 1896, p. 273—318, 3 Taf., 4 Fig.
- 1903 WILDER, H. H., The skeletal system of *Necturus maculatus* RAFINESQUE. *Mem. Boston Soc. Nat. Hist.*, Vol. 5, 1903, p. 387—439, 6 pl., 1 fig.
- 1909 WILDER, INEZ W., The lateral nasal gland in *Amphiuma*. *Journ. Morphol.*, Vol. 20, 1909, 1 pl., 7 fig.
- 1898 WINSLOW, G. M., The chondrocranium of the Ichthyopsida. *Bull. Essex Inst.*, Vol. 28, 1898, p. 87—110, 1 pl.
- 1904 WINSLOW, G. M., Three cases of abnormality in Urodeles. *Tufts Coll. Stud.*, 1904, No. 8, p. 387—410, 2 pl.
- 1854 WYMAN, JEFFRIES, Structure of the heart and physiology of respiration in the *Menobranchus* and *Batrachia*s. *Proc. Boston Soc. Nat. Hist.*, Vol. 5, 1854, p. 51—52.

- 1888 ZELLER, E., Ueber die Larve des *Proteus anguineus*. Zool. Anz., Bd. 11, 1888, p. 570—572.  
 1889 ZELLER, E., Ueber die Fortpflanzung von *Proteus anguineus* und seine Larve. Jahreshefte d. Vereins f. vaterl. Naturk. in Württemberg, 1889.  
 1890 ZELLER, E., Ueber die Befruchtung der Urodelen. Zeitschr. wiss. Zool., Bd. 49, 1890, p. 583—601, 3 Fig.  
 1895 ZUCKERKANDL, E., Zur Anatomie und Entwicklungsgeschichte der Arterien des Unterschenkels und des Fußes. Anat. Hefte, Bd. 5, 1895, p. 207—291, 6 Taf.  
 1895a ZUCKERKANDL, E., Zur Anatomie und Entwicklungsgeschichte der Arterien des Vorderarmes. Anat. Hefte, Bd. 5, 1895, p. 157—205.  
 1897 ZWICK, WILLI, Beiträge zur Kenntnis des Baues und der Entwicklung der Amphibiengliedmaßen, besonders von Carpus und Tarsus. Zeitschr. wiss. Zool., Bd. 63, 1897, p. 62—114, 4 Taf.

## B. Alphabetical arrangement of authors under topics.

### 1. General Works.

#### a) Systematic.

Baird 49.  
 Barnes 26, 27.  
 Barton 07, 08, 12, 12a.  
 Beauvois 99.  
 Boulenger 82.  
 Camper 03.  
 Chapman 79, 93.  
 Cope 85, 86, 89, 89a.  
 Cuvier 00, 27, 31.  
 Eckel and Paulmire 02.  
 Erber 64.  
 Fitzinger 26, 50.  
 Freyer 46.  
 Garnier 88.  
 Gibbes 50, 53.  
 Gray 25, 57, 73.  
 Hallowell 56, 58.  
 Harlan 27, 35, 67c.  
 Harting 71.  
 Hay 92.  
 Van der Hoeven 38b, 39, 66a, 66b.  
 Holbrook 36.  
 Kingsbury 01.  
 Kingsley 02.  
 Kneeland 56.  
 Lacepede 07, 07a.  
 Le Conte 26, 28.  
 Leuckart 40.  
 Merrem 20.  
 Rafinesque 19.  
 Rein 84.  
 Rein and von Roretz 76.  
 Schlegel 38.  
 Siebold 33.  
 Stejneger 92, 96, 07, 07a.  
 Strauch 70.  
 Temminck and Schlegel 38.  
 Tschudi 37, 38.  
 Wagler 30.  
 Waite 07.  
 Wiel-Neuwied 65.

#### b) Zoological.

Abbott 73.  
 Andres 96.  
 Beale 78.  
 Beddard 03, 04.  
 Bell 35.  
 Bender 07.  
 Berry 98.  
 Bettziech 60.  
 Bjeletzki 89.  
 Blanchard 71, 71a, 91.  
 Boie 26.  
 Bonaparte 32, 38, 39.  
 Bugnion 74.  
 Chiaje 40.  
 Claus 82.  
 Configliachi and Rusconi 21.  
 Cope 95.  
 Dalton 53.  
 Davison 94, 95, 96.  
 De Kay 42.  
 Dowler 59.  
 Dumeril et Bibron 34.  
 Ecker 59.  
 Ehrenberg 62, 67, 68, 70, 72, 74.  
 Eigenmann 99.  
 Eismann 75.  
 Ellis 66.  
 Emerson 05.  
 Emery 97.  
 Erber 63, 77.  
 Eycleshymer 06.  
 Fischer, J. G. 64.  
 Fischer, Sigwart 00.  
 Frauenfeld 61.  
 Frear 82.  
 Fülleborn 94.  
 Gadow 01.  
 Geerts 83, 84.  
 Gegenbaur 98.  
 Geyer 05.  
 Gmelin 06.  
 Grote 76, 76a.  
 Grube 66.  
 Guerne 96.

Hargitt 92.  
 Harlan 23, 24, 24a, 26, 26a.  
 Hay 88.  
 Heidenhain 07.  
 Hertwig 06.  
 Van der Hoeven 38, 38a, 67, 67a, 67b.  
 Hoffmann 77.  
 Humboldt 05.  
 Hunter 66.  
 Huxley 74, 75.  
 Hyrtl 50, 65, 65a.  
 Ishikawa 00, 02, 04, 08.  
 James 23.  
 Kammerer 05, 05a.  
 Kingsley 92.  
 Kneeland 58.  
 Kölliker 89.  
 Kornhuber 64.  
 Lankes 06.  
 Le Conte 49.  
 Leuckart 21.  
 Lewis 04.  
 Leydig 53.  
 Linnaeus 66.  
 Lloyd 66.  
 Marchesetti 75.  
 Marenzeller 78.  
 Marshall 93.  
 Mayer 35.  
 Pompe van Meerdervoort 59.  
 Mettenheimer 70.  
 Meuron 93.  
 Michahelles 29, 31.  
 Milde 62.  
 Milner 74.  
 Minot 92.  
 Mitchill 22, 24, 24a.  
 Moesch 87.  
 Montgomery 80.  
 Moodie 08.  
 Morgan 03.  
 Müller, J. C. 32.  
 Müller, K. 64.  
 Neill, 28, 32.  
 Nobbe 64.

Nussbaum 07.  
 Oesterdam 89.  
 Oken 17, 36.  
 Oppel 90, 96.  
 Osawa 99, 02.  
 Osborn 81.  
 Owen 66.  
 Phisalix 97, 97a.  
 Rabl 89.  
 Reese 06.  
 Rejsek 97.  
 Rudolphi 17.  
 Rusconi 27, 28, 37, 43, 43a.  
 Ryder 90.  
 Rusconi and Contigliachi 18.  
 Sasaki 87.  
 Schlegel 37.  
 Schmidl 50.  
 Schmidt, Goddard en Van der Hoeven 64.  
 Schneider, J. G. 79.  
 Schneider, O. 73.  
 Schneider, R. 90.  
 Schreibers 01, 18, 20.  
 Siebold 28.  
 Smith, H. M. 98.  
 Smith, J. 32.  
 Smith, F. A. 28.  
 Spicer 69.  
 Stöhr 06.  
 Stricker 72.  
 Strube 88.  
 Tommasini 75.  
 Townsend 82.  
 Treviranus 20.  
 Vaillant 63, 63a.  
 Valentin 37.  
 Viator 37.  
 Wagler 28.  
 Wagner, F. 80.  
 Wagner, R. 37.  
 Weinzettl 81.  
 Weismann and Wiedersheim 78.  
 Whitman 85, 99.  
 Wiedersheim 77, 90, 99.  
 Wilder, B. G. 74, 83.  
 Wilder, H. H. 91.  
 Zeller, 88.

### c) Embryological.

Broili 03.  
 Chauvin 83, 83a.  
 Cope 86.  
 Eycleshymer 02, 04, 04a.  
 Fülleborn 94.  
 Hay 88.  
 Kammerer 07.  
 Keibel 01, 06.

Kerbert 03, 04, 05.  
 Krausz 61.  
 Mc Gregor 96.  
 Morgan 97.  
 Platt 91, 96.  
 Rabl 98.  
 Schoenichen 01.  
 Schulze, F. E. 76.  
 Schulze, O. 05.  
 Smith, B. G. 06.  
 Virchow 91.  
 Viré 04.  
 Whitman 88, 88a.  
 Wiedersheim 77, 90.  
 Zeller 89, 90.

### 2. Earliest Stages.

Andrews 98, 98a.  
 Bussy 04, 05.  
 Camerano 83.  
 Eycleshymer 95, 02.  
 Ishikawa 03, 05.  
 Kerbert 03.  
 Lange 06, 07.  
 Schmidt 04.  
 Sobotta 97.  
 Whitman and Eycleshymer 96.  
 Zeller 90.

### 3. Skeleton.

Albrecht 78.  
 Baur 85, 88, 89, 91.  
 Buchs 02.  
 Carlson 86.  
 Cope 89.  
 Davison 97.  
 Dugès 34.  
 Field 95.  
 Gadow 95.  
 Gage 82.  
 Gaupp 04, 06.  
 Gegenbaur 62.  
 Göppert 95, 96.  
 Hay 89, 90.  
 Hepburn 07.  
 Van der Hoeven 66.  
 Houghton 03.  
 Howes and Davies 88.  
 Lucas 86.  
 Meyerhofer 09.  
 Mivart 70.  
 Murray 97.  
 Nussbaum 08.  
 Parker 68, 77, 96.  
 Peter 94, 98.  
 Platt 97.  
 Reynolds 97.

Ryder 79.  
 Schauinsland 06.  
 Sieglbauer 01.  
 Smallwood 08.  
 Smith, F. 00.  
 Terry 01.  
 Van Pée 03, 03a, 04.  
 Walter 87.  
 Whipple 06.  
 Wiedersheim 77, 92.  
 Wilder, H. H. 03.  
 Winslow 98.  
 Zwick 97.

### 4. Muscles.

Braus 07.  
 Davison 94.  
 Drüner 03.  
 Dugès 34.  
 Eycleshymer 02a, 04.  
 Fürbringer 06.  
 Gage, S. P. 90.  
 Göppert 94.  
 Humphry 72.  
 Macallum 87a.  
 Maurer 06.  
 Mivart 69, 69a.  
 Platt 97.  
 Ribbing 07.  
 Schaper 02.

### 5. Blood and Vascular System.

Boas 81.  
 Claypole 93, 96.  
 Crisp 60.  
 Eisen 99.  
 Favaro 06.  
 Gage, S. H. 85a.  
 Gulliver 73.  
 Harting 58.  
 Hochstetter 06.  
 Van der Hoeven 41, 67a.  
 Joseph 05.  
 Kneeland 68.  
 Knoll 96.  
 Königstein 03.  
 Kunitomo 03.  
 Langerhans 73.  
 Macallum 92.  
 Mandl 39, 39a.  
 Marciniowski 06.  
 Miller 00a.  
 Minot 98, 00.  
 Owen 35, 42.  
 Romeisen 05.  
 Rooy 07.  
 Schmidt, H. D. 78.  
 Sobotta 02.

Vaillant 62.  
 Welikij 90.  
 Wyman 54.  
 Zuckerkandl 95, 95a.

### 6. Respiratory System.

Böttger 76.  
 Bridge 00.  
 Camerano 83a.  
 Clemens 95.  
 Drüner 01.  
 Eycleshymer 06a.  
 Göppert 98.  
 Kneeland 56.  
 Königstein 03a.  
 Maurer 88, 88a.  
 Miller 00b, 02, 05.  
 Wiedersheim 77b.  
 Wilder, H. H. 96.  
 Wyman 54.

### 7. Digestive System.

Becker 02.  
 Bensley 00.  
 Blyer 74.  
 Bolau 99.  
 Broman 04.  
 Choronshtitzky 97, 00.  
 Gage-Gage 90.  
 Gage, S. H. 85.  
 Göppert 91, 06.  
 Green 96, 97.  
 Holmgren 04.  
 Kingsbury 94.  
 Leydig 98.  
 Macallum 87.  
 Mc Gill 08a.  
 Mathews 99.  
 Meckel 19.  
 Miller 06.  
 Monti 98.  
 Nemiloff 02.  
 Oppel 95.  
 Platt 96.  
 Prenant 04.  
 Reese 04b.  
 Richter 02.  
 Riddle 09.  
 Saint-Hilaire 03.  
 Schulz 00.

### 8. Nervous System.

Allis 03.  
 Anderson 92.  
 Bender 07a.  
 Brachet 07.  
 Bugnion 74.  
 Carlson 06.

Deen 34.  
 Edinger 04.  
 Fischer, J. G. 43.  
 Galeotti 97.  
 Herrick 93, 94, 09.  
 Hirsch-Tabor 09.  
 Hoffmann 02.  
 Humphry 72.  
 Jelliffe 97.  
 Johnston, J. B. 03, 06.  
 Kingsbury 95a.  
 Klaussner 83.  
 Kühne 86.  
 Kupffer 06.  
 Levi 97.  
 Leydig 76a.  
 Macallum 87.  
 Mc Gill 08, 08a.  
 McGregor 96.  
 Miller 00.  
 Monti 99.  
 Nakagawa 91.  
 Nemiloff 02.  
 Neumayer 06.  
 Norris 05, 08, 09.  
 Osborn 83, 84, 86, 86a, 88.  
 Platt 96a.  
 Rubaschkin 01, 03.  
 Ruge 97.  
 Smirnow 90.  
 Strong 92, 95.  
 Tuerckheim 03.  
 Vaillant 62.  
 Warren 05.

### 9. Sense Organs.

Bawden 94.  
 Beer 99.  
 Bugnion 73.  
 Dubois 90.  
 Eigenmann 00, 00a.  
 Eigenmann and Denny 00, 00a.  
 Eycleshymer 93.  
 Fleissig 09.  
 Hess 89.  
 Howard 08.  
 Kingsbury 95, 95b, 03, 08.  
 Kingsley 00.  
 Kingsley and Ruddick 99.  
 Kohl 89, 91.  
 Kuhn 79.  
 Lauber 02.  
 Leydig 68, 76b.  
 Loeb 05.  
 Malbranc 75, 76.  
 Mall 93.  
 Mihalkovics 98.

Norris 94.  
 Okajima 06.  
 Okasima 06.  
 Rabl 98.  
 Reese 05a, 06a.  
 Rejsek 97a.  
 Retzius 81, 04, 05.  
 Schlamp 91, 92.  
 Schulze, F. E. 61, 70.  
 Seydel 95.  
 Studnička 97.  
 Takahashi 09.  
 Terry 06.  
 Wilder, H. H. 92.  
 Wilder, J. W. 09.

### 10. Integument.

Ancel 01.  
 Chauvin 84.  
 Eycleshymer 06b.  
 Grote 76.  
 Heidenhain 93.  
 Leydig 73, 76, 76a.  
 Merkel 80.  
 Niclogu 93.  
 Rabl 95.  
 Reese 04b, 05.  
 Schuberg 93.  
 Studer 74.

### 11. Genito-urinary Organs.

Bidder 46.  
 Broman 01.  
 Drzewina 03.  
 Duvernoy 47.  
 Felix and Bühler 06.  
 Field 94.  
 Fürbringer 77.  
 Hall 04.  
 Heidenhain 00.  
 Johnston, C. 76.  
 Joseph 05a.  
 Kampen 03.  
 Kingsbury 95.  
 Knappe 86.  
 McGregor 99.  
 Meyer 75.  
 Minot 98.  
 Piersol 89.  
 Rathke 25.  
 Reese 04, 04a.  
 Valentin 41.

### 12. Variations.

Bumpus 96.  
 Howes 93.  
 Waite 97.  
 Winslow 04.



















